

=> d que stat 146

L4 139126 SEA FILE=HCAPLUS ABB=ON PLU=ON GAMMA RAY+PFT,NT/CT  
L5 413 SEA FILE=HCAPLUS ABB=ON PLU=ON GAMMA RAY STERILIZATION+PFT/CT  
  
L6 15042 SEA FILE=HCAPLUS ABB=ON PLU=ON "STERILIZATION AND DISINFECTIO  
N"+PFT,NT/CT  
L7 6208 SEA FILE=HCAPLUS ABB=ON PLU=ON (L4 OR L OR GAMMA) AND  
(L6 OR STERIL? OR DISINFEC?)  
L8 6208 SEA FILE=HCAPLUS ABB=ON PLU=ON L7 OR L5  
L44 1 SEA FILE=REGISTRY ABB=ON PLU=ON MORPHOLINOPROPANESULFONIC  
ACID/CN  
L45 524 SEA FILE=HCAPLUS ABB=ON PLU=ON L44  
L46 1 SEA FILE=HCAPLUS ABB=ON PLU=ON L8 AND L45

=> s 117 or 128 or 132 or 134 or 140 or 143 or 146

L47 25 L17 OR L28 OR L32 OR L34 OR L40 OR L43 OR L46

=> d 147 ibib abs hitind hitstr 1-25

L47 ANSWER 1 OF 25 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2004:533527 HCAPLUS

DOCUMENT NUMBER: 141:59810

TITLE: Process for preparing a chemically modified  
fibrin-fibrillar protein (FFP) composite sheet

INVENTOR(S): Noorjahan, Sheik Eusuff; Ranganayaki, Mandyam  
Devasikamani; Radhakrishnan, Ganga; Das, Bhabendra  
Nath; Venkateswarlu, Ummadisetty; Rose, Chellan;  
Sastry, Thotapalli Parvathaleswara

PATENT ASSIGNEE(S): India

SOURCE: U.S. Pat. Appl. Publ., 6 pp.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2004124564	A1	20040701	US 2002-330477	20021230
PRIORITY APPLN. INFO.:			US 2002-330477	20021230

AB The present invention relates to a process for the preparation of a novel  
chemical

modified fibrin-fibrillar protein (FFP) composite sheet for medical  
application and the FFP composite prepared thereby. The FFP sheet finds  
potential use as a dressing aid in the treatment of various external  
wounds of different nature, which include cut wounds, burn wounds and even  
ulcers in animals and human beings.

IC ICM B29C039-02

ICS B29C071-04

INCL 264488000; 264299000

CC 63-7 (Pharmaceuticals)

IT Bleaching agents

Crosslinking agents

Gamma ray


Human

Plasticizers

Sterilization and Disinfection

(process for preparing a chemical modified fibrin-fibrillar protein (FFP))

composite sheet)  
 IT 127-08-2, Potassium acetate 127-09-3, Sodium acetate 7681-57-4  
 , Sodium metabisulfite 7727-21-1, Potassium persulfate 7727-54-0,  
 Ammonium persulfate  
 RL: CAT (Catalyst use); USES (Uses)  
 (process for preparing a chemical modified fibrin-fibrillar protein (FFP)  
 composite sheet)  
 IT 7681-57-4, Sodium metabisulfite  
 RL: CAT (Catalyst use); USES (Uses)  
 (process for preparing a chemical modified fibrin-fibrillar protein (FFP)  
 composite sheet)  
 RN 7681-57-4 HCAPLUS  
 CN Disulfurous acid, disodium salt (9CI) (CA INDEX NAME)



●2 Na

L47 ANSWER 2 OF 25 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2004:138606 HCAPLUS

DOCUMENT NUMBER: 140:160137

TITLE: Gamma-sterilizable casein  
 -soy-peptone-agar culture medium for the  
 detection of microorganisms in hydrogen  
 peroxide-containing air and on surfaces with  
 hydrogen peroxide

INVENTOR(S): Horn, Juergen  
 PATENT ASSIGNEE(S): Biotest AG, Germany  
 SOURCE: Ger. Offen., 7 pp.

CODEN: GWXXBX

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 10233346	A1	20040219	DE 2002-10233346	20020723
US 2004106186	A1	20040603	US 2003-623241	20030718 <--
EP 1394264	A1	20040303	EP 2003-16728	20030722
EP 1394264	B1	20041103		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
AT 281531	E	20041115	AT 2003-16728	20030722
ES 2230526	T3	20050501	ES 2003-3016728	20030722
HK 1065568	A1	20050715	HK 2004-106678	20040903
PRIORITY APPLN. INFO.:			DE 2002-10233346	A 20020723

AB The invention concerns a culture medium that is gamma-sterilizable and also resists the inhibiting effect of hydrogen peroxide during culturing of microorganisms; the culture medium includes 2-10% sodium thioglycolate, 5-20% sodium thiosulfate and 10-30% sodium disulfite for neutralizing hydrogen

peroxide; the effect is increased in the presence of sodium pyruvate. To protect the color indicators during gamma radiation, polyvinylpyrrolidone and MOPS are added. Thus a medium contained in a 1 L volume with water (g): Microbial Content Test Agar 23; agar containing casein, soy peptone, sodium chloride, lecithin and sorbitan monooleate 12; polyvinylpyrrolidone 10; betaine 0.03; glycine 0.05; L-cystine 0.025; L-proline 0.025; sodium pyruvate 0.25; L-asparagine 0.025; D-glucose 2.5; sodium thioglycolate 1.0; sodium disulfite 2.5; sodium thiosulfate 6.0; bromcresol purple 0.025; bromthymol blue 0.025. The mixture was autoclaved; after cooling the following sterile filtrated ingredients were added (mL): yeast extract (from a mixture of 10 g yeast in 100 mL water) 2.5; 1M phosphate buffer pH 7.3 20; 4M MOPS buffer pH 7.4 6; L-ascorbic acid (from a solution of 1 g sodium ascorbate in 2 mL water) 0.5.

- IC ICM C12Q001-04
- CC 9-11 (Biochemical Methods)
- Section cross-reference(s): 59
- ST culture medium gamma sterilization quality control
- IT antimicrobial hydrogen peroxide
- IT Acid-base indicators
- Antimicrobial agents
- Culture media
- Microorganism
- Quality control
  - Sterilization and Disinfection
  - (gamma-sterilizable casein-soy-peptone-agar culture medium for the detection of microorganisms in hydrogen peroxide-containing air and on surfaces with hydrogen peroxide)
- IT Betaines
  - RL: BSU (Biological study, unclassified); BIOL (Biological study)
  - (gamma-sterilizable casein-soy-peptone-agar culture medium for the detection of microorganisms in hydrogen peroxide-containing air and on surfaces with hydrogen peroxide)
- IT Gamma ray
  - (irradiation; gamma-sterilizable casein-soy-peptone-agar culture medium for the detection of microorganisms in hydrogen peroxide-containing air and on surfaces with hydrogen peroxide)
- IT Air analysis
  - (microorganisms; gamma-sterilizable casein-soy-peptone-agar culture medium for the detection of microorganisms in hydrogen peroxide-containing air and on surfaces with hydrogen peroxide)
- IT Sterilization and Disinfection
  - (radiation-induced,  $\gamma$ -irradiation; gamma-sterilizable casein-soy-peptone-agar culture medium for the detection of microorganisms in hydrogen peroxide-containing air and on surfaces with hydrogen peroxide)
- IT 14265-44-2, Phosphate, biological studies
  - RL: BSU (Biological study, unclassified); BIOL (Biological study)
  - (buffer; gamma-sterilizable casein-soy-peptone-agar culture medium for the detection of microorganisms in hydrogen peroxide-containing air and on surfaces with hydrogen peroxide)
- IT 76-59-5, Bromthymol blue 115-40-2, Bromcresol purple
  - RL: ARG (Analytical reagent use); BSU (Biological study, unclassified);

ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (gamma-sterilizable casein-soy-peptone-  
 agar culture medium for the detection of microorganisms in  
 hydrogen peroxide-containing air and on surfaces with  
 hydrogen peroxide)

IT 56-40-6, Glycine, biological studies 56-89-3, L-Cystine, biological  
 studies 70-47-3, L-Asparagine, biological studies 113-24-6,  
 Sodium pyruvate 147-85-3, L-Proline, biological studies 367-51-1  
 , Sodium thioglycolate 1132-61-2, MOPS 7681-57-4  
 7772-98-7, Sodium thiosulfate 9003-39-8,  
 Polyvinylpyrrolidone

RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (gamma-sterilizable casein-soy-peptone-  
 agar culture medium for the detection of microorganisms in  
 hydrogen peroxide-containing air and on surfaces with  
 hydrogen peroxide)

IT 7722-84-1, Hydrogen peroxide, biological  
 studies

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
 (Uses)

(gamma-sterilizable casein-soy-peptone-  
 agar culture medium for the detection of microorganisms in  
 hydrogen peroxide-containing air and on surfaces with  
 hydrogen peroxide)



IT 76-59-5, Bromthymol blue

RL: ARG (Analytical reagent use); BSU (Biological study, unclassified);  
 ANST (Analytical study); BIOL (Biological study); USES (Uses)

(gamma-sterilizable casein-soy-peptone-  
 agar culture medium for the detection of microorganisms in  
 hydrogen peroxide-containing air and on surfaces with  
 hydrogen peroxide)

RN 76-59-5 HCAPLUS

CN Phenol, 4,4'-(1,1-dioxido-3H-2,1-benzoxathiol-3-ylidene)bis[2-bromo-3-  
 methyl-6-(1-methylethyl)- (9CI) (CA INDEX NAME)




IT 113-24-6, Sodium pyruvate 367-51-1, Sodium thioglycolate

1132-61-2, MOPS 7681-57-4 7772-98-7, Sodium  
thiosulfate 9003-39-8, Polyvinylpyrrolidone  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(gamma-sterilizable casein-soy-peptone-  
agar culture medium for the detection of microorganisms in  
hydrogen peroxide-containing air and on surfaces with  
hydrogen peroxide)

RN 113-24-6 HCAPLUS


CN Propanoic acid, 2-oxo-, sodium salt (9CI) (CA INDEX NAME)



● Na

RN 367-51-1 HCAPLUS


CN Acetic acid, mercapto-, monosodium salt (8CI, 9CI) (CA INDEX NAME)



● Na


RN 1132-61-2 HCAPLUS

CN 4-Morpholinepropanesulfonic acid (7CI, 8CI, 9CI) (CA INDEX NAME)



RN 7681-57-4 HCAPLUS


CN Disulfurous acid, disodium salt (9CI) (CA INDEX NAME)



●2 Na

RN 7772-98-7 HCAPLUS

CN Thiosulfuric acid (H<sub>2</sub>S<sub>2</sub>O<sub>3</sub>), disodium salt (9CI) (CA INDEX NAME)



●2 Na


RN 9003-39-8 HCAPLUS

CN 2-Pyrrolidinone, 1-ethenyl-, homopolymer (9CI) (CA INDEX NAME)

CM 1

CRN 88-12-0

CMF C6 H9 N O



IT 7722-84-1, Hydrogen peroxide, biological studies

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(gamma-sterilizable casein-soy-peptone-agar culture medium for the detection of microorganisms in hydrogen peroxide-containing air and on surfaces with hydrogen peroxide)

RN 7722-84-1 HCAPLUS

CN Hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) (9CI) (CA INDEX NAME)

HO-OH

L47 ANSWER 3 OF 25 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2003:905843 HCAPLUS

DOCUMENT NUMBER: 141:6042

TITLE: Combined effects of ionizing-irradiation and different environments on Clostridium botulinum type E spores

AUTHOR(S): Lim, Y. H.; Hamdy, M. K.; Toledo, R. T.

CORPORATE SOURCE: Department of Food Science and Technology, University of Georgia, Athens, GA, 30602, USA

SOURCE: International Journal of Food Microbiology (2003), 89(2-3), 251-263

CODEN: IJFMDD; ISSN: 0168-1605

PUBLISHER: Elsevier Science Ltd.

DOCUMENT TYPE: Journal

## LANGUAGE:

English

AB We examined the combined effects of .gamma.-radiation (24 °C) on spores of Clostridium botulinum-type Eklund strain suspended in different gas-saturated Na-phosphate buffers in the absence or presence of protectors or sensitizers. Response surface methodol. (RSM) was also used to ascertain the effects of radiation on the recovery of spores using a medium containing various levels of NaCl or Na-thioglycolate. The former (<0.5%) decreased viable spore counts, but the latter (0.15%) did not. Irradiation inactivation of Eklund spores was most effective in air-saturated buffers compared to N2O and N2 gas. The Na2-EDTA (0.01 M) was the most efficient radioprotector of spores due to its reactivity toward hydroxy radicals, followed by t-butanol (0.1 M) in NO2 or N2-saturated buffers, resp. Catalase (10.0 mg ml-1) and dl-cysteine (0.1 mM) sensitized the spores during irradiated N2O or N2-saturated buffers, and NaCl (0.01 M) only sensitized spores in N2 environment. Spores frozen at -75°C for 30 days and thawed prior to use were more sensitive to radiation damage compared to freshly prepared spores. Glycerol (15%), in Na-phosphate buffer (pH 7.0, 0.06 M), protected Eklund spores and increased the number of spores from 106 to 1011 colony forming unit (CFU) ml-1, and enhanced their radiosensitivities. Seven strains of C. botulinum type E were screened for plasmids and strain BL764 had two plasmids (15.8 and 46.8 mDa), BL4028 also had two (4.4 and 13.2 mDa), BL4850 contained only one (4.9 mDa), whereas EQA, BL211, Eklund, and Beluga had none. .gamma.-Radiation (10 kGy, absorbed dose) cured the 15.8-mDa plasmid in strain BL764, but its absence yielded no changes in toxigenicity.

CC 17-5 (Food and Feed Chemistry)

ST Clostridium spore gamma irradiation


IT Clostridium botulinum  
Gamma ray sterilization  
(combined effects of ionizing-irradiation and different environments on Clostridium botulinum type E spores)

IT 60-00-4, EDTA, uses 75-65-0, t-Butanol, uses 367-51-1, Sodium thioglycolate 3374-22-9, Cysteine 7647-14-5, Sodium chloride, uses 9001-05-2, Catalase  
RL: MOA (Modifier or additive use); USES (Uses)  
(combined effects of ionizing-irradiation and different environments on Clostridium botulinum type E spores)

IT 367-51-1, Sodium thioglycolate  
RL: MOA (Modifier or additive use); USES (Uses)  
(combined effects of ionizing-irradiation and different environments on Clostridium botulinum type E spores)

RN 367-51-1 HCAPLUS

CN Acetic acid, mercapto-, monosodium salt (8CI, 9CI) (CA INDEX NAME)



● Na

REFERENCE COUNT: 44 THERE ARE 44 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L47 ANSWER 4 OF 25 HCAPLUS COPYRIGHT 2006 ACS on STN  
ACCESSION NUMBER: 2003:235486 HCAPLUS

DOCUMENT NUMBER: 138:260448  
 TITLE: Antibiotic formulations containing  
 N-methyl-2-pyrrolidone  
 INVENTOR(S): Mihalik, Richard; Carpenter, John R.; Faris, Heidi M.  
 P.  
 PATENT ASSIGNEE(S): Phoenix Scientific, Inc., USA  
 SOURCE: U.S., 4 pp.  
 CODEN: USXXAM  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 6537985	B1	20030325	US 2001-997978	20011130
CA 2468679	AA	20030612	CA 2002-2468679	20021127
WO 2003047590	A1	20030612	WO 2002-US38007	20021127
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
EP 1458398	A1	20040922	EP 2002-782387	20021127
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
PRIORITY APPLN. INFO.:			US 2001-997978	A 20011130
			WO 2002-US38007	W 20021127

AB An antibiotic formulation in a true solution is provided. This formulation contains an antibiotic and N-methyl-2-pyrrolidone. It also may include a preservative, an antioxidant, and/or an additive. The antibiotic is a  $\beta$ -lactam, such as a penicillin, a cephalosporin, other  $\beta$ -lactams, or combinations thereof. The formulation is made by dissolving the antibiotic in N-methyl-2-pyrrolidone. The antibiotic formulation is suitable for use at temps. below 0° and without agitation. Further, the antibiotic formulation in true solution can be made with non-sterile ingredients and can be filtered to remove impurities. N-methyl-2-pyrrolidone (330 mL) was warmed to 35-40° and 125 g amoxicillin in powder form was added. The mixture was stirred until most of the material was dissolved resulting in a lemon-yellow solution. The mixture was cooled to 30° and after cooling the mixture to 30°, approx. 15 g benzyl alc. (preservative) was added to the mixture. The temperature was then increased up to 50° to dissolve the antibiotic completely.

IC ICM A61K031-545

INCL 514200000; 514198000; 514199000

CC 63-6 (Pharmaceuticals)


IT 50-81-7, Vitamin C, biological studies 58-95-7, Vitamin E acetate 68-19-9, Vitamin B12 139-33-3, Edetate disodium 149-44-0, Sodium formaldehyde sulfoxylate 7681-57-4, Sodium metabisulfite

RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses) (antioxidant; antibiotic formulations containing methylpyrrolidone)

IT 57-15-8, Chlorobutanol 64-17-5, Ethyl alcohol, biological studies 65-85-0, Benzoic acid, biological studies 100-51-6, Benzyl alcohol,



biological studies 532-32-1, Sodium benzoate 2748-88-1, Myristyl  
 $\gamma$ -picolinium chloride  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (preservative; antibiotic formulations containing methylpyrrolidone)  
 IT 7681-57-4, Sodium metabisulfite  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (antioxidant; antibiotic formulations containing methylpyrrolidone)  
 RN 7681-57-4 HCAPLUS  
 CN Disulfurous acid, disodium salt (9CI) (CA INDEX NAME)



●2 Na

REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS  
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L47 ANSWER 5 OF 25 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2003:4785 HCAPLUS

DOCUMENT NUMBER: 138:61389

TITLE: Sterilization of polymeric bioactive  
 coatings for medical goods

INVENTOR(S): Timm, Debra A.; Hui, Henry K.; Roller, Mark B.;  
 Melican, Mora C.; Hossainy, Syed

PATENT ASSIGNEE(S): Ethicon, Inc., USA

SOURCE: Eur. Pat. Appl., 13 pp.

CODEN: EPXXDW

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1270018	A1	20030102	EP 2002-254563	20020628
EP 1270018	B1	20050413		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
US 2003175408	A1	20030918	US 2001-897657	20010629
US 6787179	B2	20040907		
AU 2002048812	A5	20030102	AU 2002-48812	20020617
CA 2391899	AA	20021229	CA 2002-2391899	20020627
JP 2003047645	A2	20030218	JP 2002-191227	20020628
EP 1559434	A1	20050803	EP 2005-75683	20020628
R: DE, ES, FR, GB, IT				
ES 2239701	T3	20051001	ES 2002-2254563	20020628
PRIORITY APPLN. INFO.:			US 2001-897657	A 20010629
			EP 2002-254563	A3 20020628

AB The invention provides a method for single-step surface modification,  
 grafting and sterilization for bioactive coating on materials  
 and biomaterials used in medical devices, such as catheters, tissue  
 engineering scaffolds, or drug delivery carrier materials. This may  
 include any medical device or implantable that could benefit from improved

antithrombogenic and biocompatible surfaces. Other relevant device examples may include heparin or urokinase coated stents to reduce clotting and restenosis, dental or ophthalmol. implants. These materials may be comprised of a variety of polymeric compns. such as, polyurethane, polyester, polytetrafluoroethylene, polyethylene, polymethyl methacrylate, polyHEMA, polyvinyl alc., polysiloxanes, polylactic or glycolic acids, polycaprolactone. The substrates can also be metal, ceramics or biol. derived materials. For the sterilization process, PEG incorporation (O/C) and heparin grafting (S) were higher compared to other processes. The coating solution was a 1:1 dilution of PEG Acrylate (1.9%) + heparin (2.85%) + hyaluronic acid (0.5%) in solution of 0.5% Tween H2O.

- IC ICM A61L002-14
- ICS A61L002-18; A61L002-20; A61L027-28; A61L029-08; A61L031-08; A61L033-00; A61M025-00; B05D007-24
- CC 63-7 (Pharmaceuticals)
- ST sterilization bioactive coating polymer medical
- IT Prosthetic materials and Prosthetics  
(antithrombogenic; sterilization of polymeric bioactive coatings for medical goods)
- IT Polyesters, biological studies  
RL: DEV (Device component use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(caprolactone-based; sterilization of polymeric bioactive coatings for medical goods)
- IT Medical goods  
(catheters; sterilization of polymeric bioactive coatings for medical goods)
- IT ABS rubber  
Fluoropolymers, biological studies  
Polycarbonates, biological studies  
Polyesters, biological studies  
Polyethers, biological studies  
Polysiloxanes, biological studies  
Polysulfones, biological studies  
Polyurethanes, biological studies  
RL: DEV (Device component use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(coating material; sterilization of polymeric bioactive coatings for medical goods)
- IT Phosphatidylcholines, biological studies  
Polyoxyalkylenes, biological studies  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(coating material; sterilization of polymeric bioactive coatings for medical goods)
- IT Polyamides, biological studies  
Polyesters, biological studies  
RL: DEV (Device component use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(coating materials; sterilization of polymeric bioactive coatings for medical goods)
- IT Sterilization and Disinfection  
(electron beam; sterilization of polymeric bioactive coatings for medical goods)
- IT Polysiloxanes, biological studies  
RL: DEV (Device component use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(fluorine-containing; sterilization of polymeric bioactive coatings for medical goods)
- IT Dental materials and appliances

- (implants; sterilization of polymeric bioactive coatings for medical goods)
- IT Polyesters, biological studies  
RL: DEV (Device component use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(lactic acid-based; sterilization of polymeric bioactive coatings for medical goods)
- IT Polyimides, biological studies  
Polyketones  
RL: DEV (Device component use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(polyether-; sterilization of polymeric bioactive coatings for medical goods)
- IT Polyethers, biological studies  
RL: DEV (Device component use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(polyimide-; sterilization of polymeric bioactive coatings for medical goods)
- IT Polyethers, biological studies  
RL: DEV (Device component use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(polyketone-; sterilization of polymeric bioactive coatings for medical goods)
- IT Fluoropolymers, biological studies  
RL: DEV (Device component use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(polysiloxane-; sterilization of polymeric bioactive coatings for medical goods)
- IT Bone  
Ceramics  
Coating materials  
Coral  
Drug delivery systems  
Gamma ray sterilization  
Medical goods  
Plasma  
Sterilization and Disinfection  
(sterilization of polymeric bioactive coatings for medical goods)
- IT Ethylene-propylene rubber  
Fluoropolymers, biological studies  
Glass, biological studies  
Metals, biological studies  
Plastics, biological studies  
Polymers, biological studies  
Polyoxymethylenes, biological studies  
Polyoxyphenylenes  
Synthetic rubber, biological studies  
RL: DEV (Device component use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(sterilization of polymeric bioactive coatings for medical goods)
- IT Collagens, biological studies  
Elastins  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(sterilization of polymeric bioactive coatings for medical goods)
- IT 9003-56-9  
RL: DEV (Device component use); THU (Therapeutic use); BIOL (Biological

study); USES (Uses)  
 (abs rubber, coating material; sterilization of polymeric bioactive coatings for medical goods)

IT 9002-84-0, PTFE 9002-88-4, Polyethylene 9002-89-5, Poly(vinyl alcohol) 9003-07-0, Polypropylene 9004-61-9, Hyaluronic acid 25038-59-9, PET, biological studies 25249-16-5, Poly(2-hydroxyethyl methacrylate)  
 RL: DEV (Device component use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (coating material; sterilization of polymeric bioactive coatings for medical goods)

IT 9003-11-6 9003-39-8, Polyvinylpyrrolidone 9003-53-6, Polystyrene 9005-49-6, Heparin, biological studies 9039-53-6, Urokinase 25213-24-5, Vinyl alcohol-vinyl acetate copolymer 25322-68-3, Polyethylene glycol 25322-69-4, Polypropylene glycol 25721-76-0, Polyethylene glycol dimethacrylate 25736-86-1, Polyethylene glycol methacrylate 25852-47-5, Polyethylene glycol methacrylate 26403-58-7, Polyethylene glycol monoacrylate 26570-48-9, Polyethylene glycol diacrylate 28158-16-9, Polyethylene glycol diacrylate 53123-88-9, Rapamycin  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (coating material; sterilization of polymeric bioactive coatings for medical goods)

IT 9010-79-1  
 RL: DEV (Device component use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (ethylene-propylene rubber, sterilization of polymeric bioactive coatings for medical goods)

IT 7722-84-1, Hydrogen peroxide, processes  
 RL: PEP (Physical, engineering or chemical process); PYP (Physical process); PROC (Process)  
 (sterilization by; sterilization of polymeric bioactive coatings for medical goods)


IT 1398-61-4, Chitin 7429-90-5, Aluminum, biological studies 7440-32-6, Titanium, biological studies 7631-86-9, Silica, biological studies 9002-86-2, PVC 9003-56-9, Acrylonitrile-butadiene-styrene copolymer 9011-14-7, PMMA 9016-80-2, Polymethylpentene 11114-92-4 12597-68-1, Stainless steel, biological studies 24937-79-9, PVDF 24980-41-4, Polycaprolactone 25248-42-4, Polycaprolactone 26009-03-0, Polyglycolic acid 26023-30-3, Poly[oxy(1-methyl-2-oxo-1,2-ethanediyl)] 26100-51-6, Polylactic acid 26124-68-5, Polyglycolic acid 29223-92-5, Poly(p-dioxanone) 31621-87-1, Poly(p-dioxanone), SRU 31852-84-3, Polytrimethylene carbonate 50862-75-4, Poly(oxycarbonyloxy-1,3-propanediyl) 52013-44-2, Nitinol  
 RL: DEV (Device component use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (sterilization of polymeric bioactive coatings for medical goods)

IT 9003-39-8, Polyvinylpyrrolidone  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (coating material; sterilization of polymeric bioactive coatings for medical goods)

RN 9003-39-8 HCAPLUS  
 CN 2-Pyrrolidinone, 1-ethenyl-, homopolymer (9CI) (CA INDEX NAME)

CM 1

CRN 88-12-0  
 CMF C6 H9 N O



IT 7722-84-1, Hydrogen peroxide, processes  
 RL: PEP (Physical, engineering or chemical process); PYP (Physical process); PROC (Process)  
 (sterilization by; sterilization of polymeric bioactive coatings for medical goods)  
 RN 7722-84-1 HCAPLUS  
 CN Hydrogen peroxide (H2O2) (9CI) (CA INDEX NAME)

HO-OH

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L47 ANSWER 6 OF 25 HCAPLUS COPYRIGHT 2006 ACS on STN  
 ACCESSION NUMBER: 2002:937303 HCAPLUS  
 DOCUMENT NUMBER: 138:20443  
 TITLE: Endocrine disruptor screening using DNA chips of endocrine disruptor-responsive genes  
 INVENTOR(S): Kondo, Akihiro; Takeda, Takeshi; Mizutani, Shigetoshi; Tsujimoto, Yoshimasa; Takashima, Ryokichi; Enoki, Yuki; Kato, Ikunoshin  
 PATENT ASSIGNEE(S): Takara Bio Inc., Japan  
 SOURCE: Jpn. Kokai Tokkyo Koho, 386 pp.  
 CODEN: JKXXAF  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2002355079	A2	20021210	JP 2002-69354	20020313
PRIORITY APPLN. INFO.:			JP 2001-73183	A 20010314
			JP 2001-74993	A 20010315
			JP 2001-102519	A 20010330

AB A method and kit for detecting endocrine-disrupting chems. using DNA microarrays are claimed. The method comprises preparing a nucleic acid sample containing mRNAs or cDNAs originating in cells, tissues, or organisms which have been brought into contact with a sample containing the endocrine disruptor. The nucleic acid sample is hybridized with DNA microarrays having genes affected by the endocrine disruptor or DNA fragments originating in these genes have been fixed. The results obtained are then compared with the results obtained with the control sample to select the gene affected by the endocrine disruptor. Genes whose expression is altered by tri-Bu tin, 4-octaphenol, 4-nonylphenol, di-N-Bu phthalate, dichlorohexyl phthalate, octachlorostyrene, benzophenone, diethylhexyl phthalate, diethylstilbestrol (DES), and 17- $\beta$  estradiol (E2), were found in mice by DNA chip anal.

IC ICM C12N015-09

ICS C12N015-09; C12Q001-02; C12Q001-68; G01N033-53; G01N037-00  
CC 3-1 (Biochemical Genetics)  
Section cross-reference(s): 2, 4, 5, 9, 13  
IT Proteins  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(14-3-3, 14-3-3 protein  $\gamma$  ; endocrine disruptor  
screening using DNA chips of endocrine disruptor-responsive genes)  
IT Transcription factors  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(AP-2 (activator protein 2), transcription factor AP-2  $\gamma$   
(activating enhancer binding protein 2  $\gamma$  ); endocrine  
disruptor screening using DNA chips of endocrine disruptor-responsive  
genes)  
IT Transcription factors  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(AP-2 (activator protein 2), transcription factor AP2  $\gamma$   
; endocrine disruptor screening using DNA chips of endocrine  
disruptor-responsive genes)  
IT Immunoglobulin receptors  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(Fc receptor IgE high affinity I  $\gamma$  polypeptide;  
endocrine disruptor screening using DNA chips of endocrine  
disruptor-responsive genes)  
IT G proteins (guanine nucleotide-binding proteins)  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(G protein,  $\gamma$  3 subunit; endocrine disruptor screening  
using DNA chips of endocrine disruptor-responsive genes)  
IT Proteins  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(KIAA0410; endocrine disruptor screening using DNA chips of  
endocrine disruptor-responsive genes)  
IT Proteins  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(MGP (matrix  $\gamma$  -carboxyglutamic acid-containing protein),  
matrix Gla protein; endocrine disruptor screening using DNA chips of  
endocrine disruptor-responsive genes)  
IT Retinoic acid receptors  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(RAR- $\gamma$  ; endocrine disruptor screening using DNA chips  
of endocrine disruptor-responsive genes)  
IT Proteins  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(RAS oncogene family protein RAB5A; endocrine disruptor screening using  
DNA chips of endocrine disruptor-responsive genes)  
IT Proteins  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(RSP5 protein; endocrine disruptor screening using DNA chips of  
endocrine disruptor-responsive genes)  
IT Glycoproteins  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(Rhesus blood group-associated A glycoprotein; endocrine  
disruptor screening using DNA chips of endocrine disruptor-responsive  
genes)  
IT Proteins  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(Ric, sequence homolog; endocrine disruptor  
screening using DNA chips of endocrine disruptor-responsive genes)  
IT Proteins  
RL: BSU (Biological study, unclassified); BIOL (Biological study)

- (Ring3; endocrine disruptor screening using DNA chips of endocrine disruptor-responsive genes)
- IT Calcium-binding proteins  
RL: BSU (Biological study, unclassified); BIOL (Biological study) (S-100, S100 calcium binding protein A8 (calgranulin A); endocrine disruptor screening using DNA chips of endocrine disruptor-responsive genes)
- IT Calcium-binding proteins  
RL: BSU (Biological study, unclassified); BIOL (Biological study) (S-100, S100 calcium binding protein AS (calgranulin); endocrine disruptor screening using DNA chips of endocrine disruptor-responsive genes)
- IT Proteins  
RL: BSU (Biological study, unclassified); BIOL (Biological study) (SEC61  $\gamma$  subunit; endocrine disruptor screening using DNA chips of endocrine disruptor-responsive genes)
- IT Proteins  
RL: BSU (Biological study, unclassified); BIOL (Biological study) (X11  $\gamma$  protein; endocrine disruptor screening using DNA chips of endocrine disruptor-responsive genes)
- IT Proteins  
RL: BSU (Biological study, unclassified); BIOL (Biological study) (adducin, adducin 2 ( $\beta$ ), 3 ( $\gamma$ ); endocrine disruptor screening using DNA chips of endocrine disruptor-responsive genes)
- IT Proteins  
RL: BSU (Biological study, unclassified); BIOL (Biological study) (attachment, N-ethylmaleimide-sensitive factor attachment protein  $\gamma$ ; endocrine disruptor screening using DNA chips of endocrine disruptor-responsive genes)
- IT Molecular chaperones  
RL: BSU (Biological study, unclassified); BIOL (Biological study) (chaperonin, subunit 3 ( $\gamma$ ); endocrine disruptor screening using DNA chips of endocrine disruptor-responsive genes)
- IT Initiation factors (protein formation)  
RL: BSU (Biological study, unclassified); BIOL (Biological study) (eukaryotic initiation factor 4  $\gamma$ ; endocrine disruptor screening using DNA chips of endocrine disruptor-responsive genes)
- IT Initiation factors (protein formation)  
RL: BSU (Biological study, unclassified); BIOL (Biological study) (eukaryotic translation initiation factor 4  $\gamma$ , 3; endocrine disruptor screening using DNA chips of endocrine disruptor-responsive genes)
- IT Proteins  
RL: BSU (Biological study, unclassified); BIOL (Biological study) (female sterile homeotic-related gene 1; endocrine disruptor screening using DNA chips of endocrine disruptor-responsive genes)
- IT Fibrinogens  
RL: BSU (Biological study, unclassified); BIOL (Biological study) (fibrinogen  $\gamma$  A and B chain; endocrine disruptor screening using DNA chips of endocrine disruptor-responsive genes)
- IT Fibrinogens  
RL: BSU (Biological study, unclassified); BIOL (Biological study) (fibrinogen  $\gamma$  chain; endocrine disruptor screening using DNA chips of endocrine disruptor-responsive genes)
- IT G proteins (guanine nucleotide-binding proteins)  
RL: BSU (Biological study, unclassified); BIOL (Biological study) (guanine nucleotide binding protein (6 protein)  $\gamma$  transducing activity polypeptide 2; endocrine disruptor screening using

- DNA chips of endocrine disruptor-responsive genes)
- IT Proteins  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(hydrogen peroxide inducible protein 53; endocrine  
disruptor screening using DNA chips of endocrine disruptor-responsive  
genes)
- IT Transcription factors  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(interferon dependent pos. acting, factor 3  $\gamma$  ;  
endocrine disruptor screening using DNA chips of endocrine  
disruptor-responsive genes)
- IT Transcription factors  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(interferon-stimulated transcription factor 3,  $\gamma$   
(48-kDa); endocrine disruptor screening using DNA chips of endocrine  
disruptor-responsive genes)
- IT Laminins  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(laminin  $\gamma$  1; endocrine disruptor screening using DNA  
chips of endocrine disruptor-responsive genes)
- IT Laminins  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(laminin  $\gamma$  2; endocrine disruptor screening using DNA  
chips of endocrine disruptor-responsive genes)
- IT Proteins  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(odz (odd Oz/ten-m), sequence homolog; endocrine disruptor  
screening using DNA chips of endocrine disruptor-responsive genes)
- IT Peptides, biological studies  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(oligopeptides, proton-dependent high affinity oligopeptide  
transporter PepT2; endocrine disruptor screening using DNA chips of  
endocrine disruptor-responsive genes)
- IT Actins  
Synaptophysin  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(sequence homolog; endocrine disruptor screening using DNA chips of  
endocrine disruptor-responsive genes)
- IT Ribonucleoproteins  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(snRNP (small nuclear ribonucleoprotein), UCR2 U2 small nuclear  
ribonucleoprotein auxiliary factor 35-kDa subunit related protein 2;  
endocrine disruptor screening using DNA chips of endocrine  
disruptor-responsive genes)
- IT 366806-33-9, Casein kinase II  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(casein kinase II  $\alpha$ 2 polypeptide; endocrine disruptor  
screening using DNA chips of endocrine disruptor-responsive genes)
- IT 178303-43-0, Caseinolytic proteinase X  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(caseinolytic protease X; endocrine disruptor screening using  
DNA chips of endocrine disruptor-responsive genes)
- IT 1393-25-5, Secretin 9000-94-6, Antithrombin-III 9001-05-2, Catalase  
9001-16-5, Cytochrome c oxidase 9001-40-5, Glucose-6 phosphate  
dehydrogenase 9001-42-7,  $\alpha$ -Glucosidase 9001-45-0, Glucuronidase  
 $\beta$  9001-48-3, Glutathione reductase 9001-50-7,  
Glyceraldehyde-3-phosphate dehydrogenase 9001-61-0, Cytosolic  
aminopeptidase 9001-63-2, Lysozyme 9001-64-3, Malate dehydrogenase  
9001-91-6, Plasminogen 9002-03-3, Dihydrofolate reductase 9002-62-4,



Prolactin, biological studies 9004-02-8, Lipoprotein lipase 9012-37-7,  
 Aminoacylase-1 9012-93-5, Ferrochelatase 9013-10-9,  
 Glucosamine-6-phosphate isomerase 9013-93-8, Phospholipase 9014-18-0,  
 Nicotinamide nucleotide transhydrogenase 9023-44-3, Tryptophanyl-tRNA  
 synthetase 9023-48-7, Seryl-tRNA synthetase 9023-56-7, CTP synthase  
 9023-58-9, Argininosuccinate synthetase 9023-69-2, Asparagine synthetase  
 9023-88-5 9023-90-9, Methylmalonyl-CoA mutase 9024-20-8, Ribulose  
 phosphate 3 epimerase 9024-60-6, Ornithine decarboxylase 9024-70-8,  
 Uroporphyrinogen decarboxylase 9025-15-4, Biotinidase 9025-73-4,  
 Phosphoserine phosphatase 9026-00-0, Lysosomal acid lipase 9026-23-7,  
 Carbamoyl-phosphate synthase 9026-24-8, Thiamin pyrophosphokinase  
 9026-30-6, Poly (A) polymerase 9026-67-9, Choline kinase 9027-13-8,  
 Enoyl-CoA hydratase 9027-32-1, Aspartyl-tRNA synthetase 9027-44-5,  
 Hydroxymethylglutaryl-CoA synthase 9027-67-2, Terminal deoxynucleotidyl  
 transferase 9027-80-9, Adenine phosphoribosyl transferase 9027-81-0,  
 Adenylosuccinate Lyase 9028-21-1, Sorbitol dehydrogenase 9028-40-4,  
 3-Hydroxyacyl-CoA dehydrogenase 9028-61-9, Estradiol  
 17 $\beta$ -dehydrogenase 9029-17-8, Pyrroline-5-carboxylate reductase  
 9029-38-3, Sulfite oxidase 9030-22-2, Uridine phosphorylase 9030-23-3,  
 Thymidine phosphorylase 9030-24-4, Uracil phosphoribosyltransferase  
 9030-38-0 9030-53-9, Galactokinase 9030-83-5, 3-Hydroxy-3-  
 methylglutaryl-CoA lyase 9031-19-0, Saccharopine dehydrogenase  
 9031-37-2, Ceruloplasmin 9031-71-4, Alanine-tRNA synthetase 9031-82-7,  
 Amidophosphoribosyltransferase 9031-86-1, Aspartoacylase 9032-03-5,  
 Phosphoribosylaminoimidazolecarboxamide formyltransferase 9032-25-1,  
 NADH cytochrome B5 reductase 9032-59-1, Fumarylacetoacetate hydrolase  
 9032-71-7, 2,3-Oxidosqualene-lanosterol cyclase 9032-73-9, Neuropathy  
 target esterase 9032-88-6, Fumarate hydratase 9033-27-6,  
 Isopentenyl-diphosphate  $\delta$  isomerase 9035-39-6, Cytochrome b5  
 9035-42-1, Cytochrome c1 9037-62-1, Glycyl-tRNA synthetase 9037-65-4,  
 $\alpha$ -L-Fucosidase 9040-59-9, Phosphodiesterase 1 9042-64-2, DOPA  
 decarboxylase 9045-77-6, Fatty acid synthase 9047-22-7, Cathepsin B  
 9054-54-0, Transacylase 9054-84-6, Xanthine dehydrogenase 9059-11-4,  
 Amine oxidase 9059-48-7, Sepiapterin reductase 9067-83-8,  
 Phosphatidate cytidyltransferase 9068-16-0, Poly(ADP ribose)  
 glycohydrolase 9068-41-1, Carnitine palmitoyltransferase 9074-91-3,  
 Hydroxymethylbilane synthase 9075-29-0, 3 Phosphoglycerate dehydrogenase  
 9075-78-9, Ethanolamine kinase 9075-81-4,  $\beta$ -Galactoside  
 $\alpha$ -2,6-sialyltransferase 9076-57-7, Histone deacetylase  
 9076-84-0, Coproporphyrinogen oxidase 12651-28-4, Transcobalamin 2  
 37184-63-7, Myoinositol 1-monophosphatase 37205-49-5,  
 Methylmalonate-semialdehyde dehydrogenase 37211-69-1,  
 2,3-Bisphosphoglycerate mutase 37237-43-7, Galactosyltransferase  
 $\beta$ -1,4-GalT V 37255-37-1, E.C. 1.3.3.2 37255-38-2, Glutaryl-CoA  
 dehydrogenase 37259-54-4, DTDP-glucose 4,6-dehydratase 37274-61-6,  
 Isovaleryl-CoA dehydrogenase 37277-82-0, Spermidine synthase  
 37341-57-4, Succinate:CoA ligase 39471-28-8, Deoxyguanosine kinase  
 51110-01-1, Somatostatin 52660-18-1, Casein kinase 1  
 55354-43-3, Arylsulfatase B 59088-23-2, Dihydroorotate dehydrogenase  
 59298-90-7, UDP-galactose:glucosylceramide  $\beta$ 1,4-galactosyltransferase  
 60320-99-2, N-Acetylglucosamine-6-sulfatase 65997-74-2, Cathepsin F  
 67339-00-8,  $\alpha$ 2,8-Sialyltransferase 67763-97-7, Insulin-like growth  
 factor 2 79079-11-1, Calpastatin 80295-40-5, Complement C2  
 80295-48-3, Complement C4 80295-62-1, Complement factor B 80295-65-4,  
 Complement factor H 81181-72-8,  $\gamma$  -Glutamyl carboxylase  
 81611-75-8, Fructose-2,6-bisphosphatase 82062-90-6, NAD-dependent  
 methylenetetrahydrofolate dehydrogenase 82391-38-6, Branched chain keto  
 acid dehydrogenase kinase 82707-54-8, Neprilysin 83268-44-4  
 86480-67-3, Ubiquitin carboxyl-terminal hydrolase 87683-70-3,

Pterin-4 $\alpha$ -carbinolamine dehydratase 90698-32-1, Leukotriene C4  
 synthase 101149-94-4, Tripeptidyl peptidase II 102577-19-5, Neuromedin  
 B 109136-49-4, Ubiquitin-specific protease 122320-05-2, Secretory  
 leukocyte protease inhibitor 124861-55-8 137632-08-7, Mitogen  
 activated protein kinase 1 141349-86-2, Cyclin-dependent kinase 2  
 141467-21-2, Calcium/calmodulin-dependent protein kinase I 142243-03-6,  
 Plasminogen activator inhibitor type II 142539-77-3, Mast cell protease  
 5 142805-56-9, DNA topoisomerase II 143180-75-0, DNA topoisomerase I  
 144114-16-9, Focal adhesion kinase 144697-17-6, c-Src tyrosine kinase  
 145809-21-8, Tissue inhibitor of metalloproteinase 3 146480-35-5, Matrix  
 metalloproteinase 2 146480-49-1, MMCP-6 protease 146838-30-4,  
 Mitogen-activated protein kinase-activated protein kinase 2 147014-97-9,  
 Cyclin dependent kinase 4 149316-81-4, 2-Hydroxyphytanoyl-CoA lyase  
 149371-24-4, Neurolysin 150605-50-8, Neuronal tyrosine/threonine  
 phosphatase 1 152478-57-4, Janus kinase 2 153190-47-7, Gene PTK2  
 tyrosine kinase 165245-94-3, NIMA-related kinase 2 165245-99-8,  
 Polo-like kinase 167397-96-8, Interleukin-1 receptor-associated kinase  
 169277-44-5, Sphingosine-1-phosphate phosphatase 169592-62-5,  
 Cyclin-dependent kinase 10 170780-57-1, LIM kinase 172306-41-1,  
 Protein kinase PCTAIRE-1 172399-47-2, BOMAPIN 173585-04-1,  
 Integrin-linked kinase 176023-64-6, Mitogen-activated protein kinase 12  
 178037-70-2, Serum and glucocorticoid regulated protein kinase  
 180189-96-2, Caspase 9 182372-15-2, Caspase 6 184049-62-5, Gene DUSP6  
 MAP kinase phosphatase 187247-72-9, Endonuclease G 188417-84-7,  
 Vascular endothelial growth factor C 191359-13-4, MAP kinase-interacting  
 kinase 1 192230-91-4, Mitogen-activated protein kinase kinase 4  
 194739-73-6, Mitogen-activated protein kinase kinase 6 196717-99-4,  
 Prenylcysteine lyase 206566-35-0, Molybdopterin synthase sulfurylase  
 212625-17-7, SPAK protein kinase 214210-47-6, Neuropilin 1  
 216503-96-7, Caspase 11 223610-95-5, Matrix metalloproteinase MMP-23  
 230951-53-8, Caspase 12 252852-50-9, SUMO-1 conjugate proteinase  
 252901-98-7, Tousled-like kinase 1 258336-77-5, UNC51.2 serine/threonine  
 kinase 288307-53-9, Inositol 1,3,4-trisphosphate 5/6 kinase  
 292850-69-2, Nardilysin 306298-47-5, MAP kinase phosphatase-1  
 321976-25-4, Sialyltransferase 324751-96-4, Stanniocalcin 2  
 324752-01-4, Stanniocalcin 1 327046-95-7, Mitogen activated protein  
 kinase kinase 5 335135-28-9, Cytochrome P450 2D10 338969-69-0,  
 Cytochrome P450 2F2 353498-78-9, Mitogen activated protein kinase 6  
 362516-16-3, Conserved helix-loop-helix ubiquitous kinase 374936-45-5,  
 Cytochrome P450 2C40 409105-92-6, Microtubule-associated testis-specific  
 serine/threonine protein kinase 440356-82-1, Cytochrome P450 7B1  
 443906-18-1, Receptor protein tyrosine phosphatase K 464896-43-3,  
 Transmembrane serine proteinase 475489-73-7, Calcium/calmodulin-  
 dependent protein kinase II 478187-31-4, P 450 2J6

RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (endocrine disruptor screening using DNA chips of endocrine  
 disruptor-responsive genes)

IT 172522-01-9, 5'-AMP-activated protein kinase

RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (isoform  $\gamma$  -1, subunit AAKG; endocrine disruptor  
 screening using DNA chips of endocrine disruptor-responsive genes)

IT 115926-52-8, Phosphatidylinositol 3-kinase

RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (phosphatidylinositol 3-kinase, G2 domain containing  $\gamma$   
 polypeptide; endocrine disruptor screening using DNA chips of endocrine  
 disruptor-responsive genes)

IT 362479-32-1, Protein phosphatase 1

RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (protein phosphatase 1 catalytic subunit  $\beta$   $\gamma$ )

isoform; endocrine disruptor screening using DNA chips of endocrine disruptor-responsive genes)

IT 9012-90-2

RL: BSU (Biological study, unclassified); BIOL (Biological study)  
( $\gamma$ , DNA polymerase  $\gamma$ , mitochondrial;  
endocrine disruptor screening using DNA chips of endocrine  
disruptor-responsive genes)

L47 ANSWER 7 OF 25 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2002:677247 HCAPLUS

DOCUMENT NUMBER: 138:317015

TITLE: Culture media for microbiological monitoring in  
isolator with residual hydrogen  
peroxide on surfaces and in air

AUTHOR(S): Horn, Juergen; Backes, Maria; Schepp, Eleonor-C.;  
Wenz, Petra

CORPORATE SOURCE: Biotest AG, USA

SOURCE: ESTECH 2002 Proceedings: Leading the Way in the  
Century Ahead, 48th IEST Annual Technical Meeting and  
16th ICCCS International Symposium on Contamination  
Control, Anaheim, CA, United States, Apr. 28-May 1,  
2002 (2002), 92-100. Institute of Environmental  
Sciences and Technology: Rolling Meadows, Ill.  
CODEN: 69DARZ

DOCUMENT TYPE: Conference; (computer optical disk)

LANGUAGE: English

AB Isolators after fumigating with hydrogen peroxide or  
peracetic acid followed by venting may, at the start of operations, still  
have residual hydrogen peroxide concns. between 0.3  
and 6ppm in the air and between 0.5 and 3 ppm on solid surfaces. Packaged  
microbiol. media may withstand the fumigating procedures without damaging  
the fertility of the Agar, but should be examined in the actual  
sampling process operation as well. During air sampling an accumulation  
of hydrogen peroxide in the water phase of the  
Agar occurs leading to concns. of up to over 100 ppm in standard  
Tryptic Soy Casein Digest Agar, preventing the  
subsequent growth of any microorganisms. The same accumulation occurs in  
gelatin filters with residual water content used for air sampling.  
Subsequent growth of microorganism on gelatin filters exposed to  
hydrogen peroxide containing air is also not possible.  
Surface sampling of hydrogen peroxide exposed surfaces  
leads to lower recoveries of subsequently inoculated microorganisms or no  
growth in case of anaerobe spores with standard Tryptic Soy Casein  
Digest Agar. Only suitably modified Tryptic Soy Casein  
Digest Agar or stabilized D/E Agar preps. circumvent  
this problem and allow uninhibited growth of microorganisms after exposure  
to isolator environments with actual sampling procedures. The developed  
gamma-sterilized Agar strips for RCS allow  
effective air monitoring in isolators and the corresponding Contact Slide  
D/E-gamma allow effective surface sampling in isolators, as  
demonstrated by the recovery of low inocula (< 100 cfu) of all USP test  
strains after air sampling or surface sampling in fumigated isolators.  
Even the anaerobic spore forming strain *Clostridium sporogenes* ATCC11437  
does grow well on the modified media after hydrogen  
peroxide exposure. The gamma-sterilization  
procedure at 16-25 Kgray kills 10E8 cfu at 16 Kgray thereby ensuring a  
uncontaminated product.

CC 9-11 (Biochemical Methods)

ST microorganism medium agar hydrogen peroxide

fumigation isolator

IT Sampling  
(air sampling; microorganism medium and agar with peroxide neutralizing activity and  $\gamma$ -sterilization-compatible to use in fumigated isolators)

IT Culture media  
Fumigation  
Gamma ray  
(microorganism medium and agar with peroxide neutralizing activity and  $\gamma$ -sterilization-compatible to use in fumigated isolators)

IT Air  
(sampling; microorganism medium and agar with peroxide neutralizing activity and  $\gamma$ -sterilization-compatible to use in fumigated isolators)

IT 7722-84-1, Hydrogen Peroxide, biological studies  
RL: ADV (Adverse effect, including toxicity); BIOL (Biological study)  
(microorganism medium and agar with peroxide neutralizing activity and  $\gamma$ -sterilization-compatible to use in fumigated isolators)

IT 7722-84-1, Hydrogen Peroxide, biological studies  
RL: ADV (Adverse effect, including toxicity); BIOL (Biological study)  
(microorganism medium and agar with peroxide neutralizing activity and  $\gamma$ -sterilization-compatible to use in fumigated isolators)

RN 7722-84-1 HCAPLUS

CN Hydrogen peroxide (H2O2) (9CI) (CA INDEX NAME)

HO-OH

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L47 ANSWER 8 OF 25 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2002:119531 HCAPLUS

DOCUMENT NUMBER: 136:189499

TITLE: Color indicator for glutaraldehyde concentration

INVENTOR(S): Minamitani, Tamio; Ota, Shinya; Tsubaki, Tomio

PATENT ASSIGNEE(S): Oriental Pharmaceutical & Cynthetic Chemical Co., Ltd., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 6 pp.  
CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1


PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2002048780	A2	20020215	JP 2000-272322	20000804
PRIORITY APPLN. INFO.:			JP 2000-272322	20000804

AB The indicator, which simply and rapidly judges effective concentration of glutaraldehyde (I) in its preps. useful as disinfectants for medical goods, is a mixture of sulfite salts, amine compds., and pigments in the powder or solidified form. A powder mixture of Na2SO3 210, taurine 266,

Brilliant Blue 6B 1, and lactose 23 mg was dissolved in 10 mL H<sub>2</sub>O to give indicator solution. Color of 1 mL of the solution changed from blue to green at I concentration  $\geq 1.8\%$  within 30-60 s, and remained blue at I concentration  $\leq 1.6\%$ .

IC ICM G01N031-22  
ICS G01N021-78; G01N031-00  
CC 64-3 (Pharmaceutical Analysis)  
Section cross-reference(s): 9, 63  
ST disinfectant glutaraldehyde concn indicator sulfite amine pigment; Brilliant Blue sodium sulfite taurine glutaraldehyde indicator  
IT Colorimetric indicators  
Disinfectants  
(color indicator for glutaraldehyde concentration containing sulfites, amine compds., and pigments)  
IT 56-12-2,  $\gamma$ -Aminobutyric acid, uses 56-40-6, Glycine, uses 60-32-2,  $\epsilon$ -Aminocaproic acid 107-35-7, Taurine 107-95-9,  $\beta$ -Alanine 632-68-8, Japan Red 105 7631-90-5, Sodium bisulfite 7681-57-4, Sodium pyrosulfite 7757-83-7, Sodium sulfite 10117-38-1, Potassium sulfite  
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)  
(color indicator for glutaraldehyde concentration containing sulfites, amine compds., and pigments)  
IT 7681-57-4, Sodium pyrosulfite  
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)  
(color indicator for glutaraldehyde concentration containing sulfites, amine compds., and pigments)  
RN 7681-57-4 HCAPLUS  
CN Disulfurous acid, disodium salt (9CI) (CA INDEX NAME)



●2 Na

L47 ANSWER 9 OF 25 HCAPLUS COPYRIGHT 2006 ACS on STN  
ACCESSION NUMBER: 2000:688830 HCAPLUS  
DOCUMENT NUMBER: 133:360652  
TITLE: Characterization of spores of *Bacillus subtilis* which lack dipicolinic acid  
AUTHOR(S): Paidhungat, Madan; Setlow, Barbara; Driks, Adam; Setlow, Peter  
CORPORATE SOURCE: Department of Biochemistry, University of Connecticut Health Center, Farmington, CT, 06032, USA  
SOURCE: Journal of Bacteriology (2000), 182(19), 5505-5512  
CODEN: JOBAA; ISSN: 0021-9193  
PUBLISHER: American Society for Microbiology  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB Spores of *Bacillus subtilis* with a mutation in *spoVF* cannot synthesize dipicolinic acid (DPA) and are too unstable to be purified and studied in detail. However, the spores of a strain lacking the three major germinant receptors (termed  $\Delta\text{ger3}$ ), as well as *spoVF*, can be isolated, although they spontaneously germinate much more readily than  $\Delta\text{ger3}$

spores. The  $\Delta$ ger3 spoVF spores lack DPA and have higher levels of core water than  $\Delta$ ger3 spores, although sporulation with DPA restores close to normal levels of DPA and core water to  $\Delta$ ger3 spoVF spores. The DPA-less spores have normal cortical and coat layers, as observed with an electron microscope, but their core region appears to be more hydrated than that of spores with DPA. The  $\Delta$ ger3 spoVF spores also contain minimal levels of the processed active form (termed P41) of the germination protease, GPR, a finding consistent with the known requirement for DPA and dehydration for GPR autoprocessing. However, any P41 formed in  $\Delta$ ger3 spoVF spores may be at least transiently active on one of this protease's small acid-soluble spore protein (SASP) substrates, SASP- $\gamma$ . Anal. of the resistance of wild-type,  $\Delta$ ger3, and  $\Delta$ ger3 spoVF spores to various agents led to the following conclusions: (i) DPA and core water content play no role in spore resistance to dry heat, dessication, or glutaraldehyde; (ii) an elevated core water content is associated with decreased spore resistance to wet heat, hydrogen peroxide, formaldehyde, and the iodine-based disinfectant Betadine; (iii) the absence of DPA increases spore resistance to UV radiation; and (iv) wild-type spores are more resistant than  $\Delta$ ger3 spores to Betadine and glutaraldehyde. These results are discussed in view of current models of spore resistance and spore germination.

CC 10-1 (Microbial, Algal, and Fungal Biochemistry)  
 IT 50-00-0, Formaldehyde, biological studies 111-30-8, Pentanedial  
 7722-84-1, Hydrogen peroxide, biological  
 studies 25655-41-8, Betadine;  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological  
 study, unclassified); BIOL (Biological study)  
 (resistance of spores of *Bacillus subtilis* which lack dipicolinic acid)  
 IT 7722-84-1, Hydrogen peroxide, biological  
 studies 25655-41-8, Betadine;  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological  
 study, unclassified); BIOL (Biological study)  
 (resistance of spores of *Bacillus subtilis* which lack dipicolinic acid)  
 RN 7722-84-1 HCAPLUS  
 CN Hydrogen peroxide (H2O2) (9CI) (CA INDEX NAME)

HO-OH

RN 25655-41-8 HCAPLUS  
 CN 2-Pyrrolidinone, 1-ethenyl-, homopolymer, compd. with iodine (9CI) (CA  
 INDEX NAME)

CM 1

CRN 7553-56-2

CMF I2

I-I

CM 2

CRN 9003-39-8


CMF (C6 H9 N O)x

CCI PMS

CM 3

CRN 88-12-0

CMF C6 H9 N O



REFERENCE COUNT: 43 THERE ARE 43 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L47 ANSWER 10 OF 25 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1997:594530 HCAPLUS

DOCUMENT NUMBER: 127:210404

TITLE: Method for sterilization of body fluid treatment apparatus

INVENTOR(S): Kuroda, Toru; Yabushita, Hajime

PATENT ASSIGNEE(S): Asahi Medical Co., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 5 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 09220281	A2	19970826	JP 1996-354601	19870225
JP 2649224	B2	19970903	JP 1987-40148	19870225
			JP 1987-40148	A3 19870225

PRIORITY APPLN. INFO.: JP 1987-40148 A3 19870225

AB A method for sterilization of body fluid treatment apparatus [e.g. hemodialyzer] involves: soaking in a solution of antioxidants selected from Na pyrosulfite, Na sulfite, Na bisulfite, acetone sodium bisulfite, sodium formaldehyde sulfoxylate, sodium hydrosulfite and ascorbic acid and then . gamma.-radiation treatment.

IC ICM A61M001-14  
ICS A61M001-14; A61L002-08

CC 63-7 (Pharmaceuticals)

ST sterilization body fluid treatment app; hemodialyzer  
sterilization pyrosulfite radiation; antioxidant radiation  
hemodialyzer sterilization


IT Dialyzers  
(hemodialyzers; sterilization of body fluid treatment apparatus)

IT Antioxidants  
Gamma ray  
(in sterilization of body fluid treatment apparatus)

IT Apparatus  
Body fluid  
Sterilization and Disinfection  
(sterilization of body fluid treatment apparatus)

IT 50-81-7, Ascorbic acid, biological studies 149-44-0, Sodium formaldehyde sulfoxylate 540-92-1, Acetone sodium bisulfite 7631-90-5, Sodium

bisulfite 7681-57-4, Sodium pyrosulfite 7757-83-7, Sodium sulfite 7775-14-6, Sodium hydrosulfite  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (in sterilization of body fluid treatment apparatus)  
 IT 7681-57-4, Sodium pyrosulfite  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (in sterilization of body fluid treatment apparatus)  
 RN 7681-57-4 HCAPLUS  
 CN Disulfurous acid, disodium salt (9CI) (CA INDEX NAME)



●2 Na

L47 ANSWER 11 OF 25 HCAPLUS COPYRIGHT 2006 ACS on STN  
 ACCESSION NUMBER: 1996:476855 HCAPLUS  
 DOCUMENT NUMBER: 125:123805  
 TITLE: Sunscreen-wound healing composition  
 INVENTOR(S): Martin, Alain  
 PATENT ASSIGNEE(S): Warner-Lambert Company, USA  
 SOURCE: PCT Int. Appl., 137 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 28  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9617624	A1	19960613	WO 1995-US12848	19951005
W: AU, CA, JP, MX, NZ, SG				
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
US 5674912	A	19971007	US 1995-446979	19950522
AU 9538596	A1	19960626	AU 1995-38596	19951005
AU 690366	B2	19980423		
EP 796107	A1	19970924	EP 1995-936858	19951005
EP 796107	B1	20030108		
R: BE, DE, FR, GB, IT, LU, NL				
ZA 9510376	A	19971006	ZA 1995-10376	19951206
PRIORITY APPLN. INFO.:				
			US 1994-350918	A 19941207
			US 1995-446979	A 19950522
			US 1991-663500	B1 19910301
			US 1993-53922	B2 19930426
			WO 1995-US12848	W 19951005

AB The present invention pertains to therapeutic sunscreen-wound healing compns. useful to minimize and treat sunburn damage. The compns. comprise a therapeutically effective amount of (1) a sunscreen agent; (2) an anti-inflammatory; and, (3) a wound healing composition. In one embodiment the wound healing composition comprises (a) pyruvate; (b) an antioxidant; and (c) a mixture of saturated and unsatd. fatty acids. The therapeutic sunscreen-wound healing compns. may be utilized in a wide variety of pharmaceutical products. This invention also relates to methods for preparing and using the



therapeutic sunscreen-wound healing compns. and the pharmaceutical products in which the therapeutic compns. may be used.

IC ICM A61K045-06

CC 63-7 (Pharmaceuticals)

Section cross-reference(s): 62

IT Antihistaminics

Antioxidants

Bactericides, Disinfectants, and Antiseptics

Fungicides and Fungistats

Immunostimulants

Inflammation inhibitors

Nutrients

Sunburn and Suntan

Sunscreens

Virucides and Virustats

Wound healing promoters

(sunscreen-wound healing compns. for treatment of sunburn)

IT 50-02-2, Dexamethasone 50-21-5, Lactic acid, biological studies  
50-23-7, Hydrocortisone 50-24-8, Prednisolone 50-78-2, Aspirin  
50-81-7, Ascorbic acid, biological studies 53-03-2, Prednisone  
53-06-5, Cortisone 53-86-1, Indomethacin 57-10-3, Hexadecanoic acid,  
biological studies 57-11-4, Octadecanoic acid, biological studies  
57-13-6, Urea, biological studies 58-95-7, Vitamin E acetate 59-02-9,  
 $\alpha$ -Tocopherol 60-33-3, 9,12-Octadecadienoic acid (Z,Z)-, biological  
studies 61-68-7, Mefenamic acid 68-26-8, Retinol 76-25-5,  
Triamcinolone acetone 79-80-1, 3,4-Didehydroretinol 89-57-6,  
Mesalamine 112-80-1, 9-Octadecenoic acid (Z)-, biological studies  
113-24-6, Sodium pyruvate 118-60-5, 2-Ethylhexyl salicylate  
119-13-1,  $\delta$ -Tocopherol 124-94-7, Triamcinolone 127-17-3, Pyruvic  
acid, biological studies 127-17-3D, Pyruvic acid, Manganese complexes  
131-57-7, Oxybenzone 134-09-8, Menthyl anthranilate 143-07-7,  
Dodecanoic acid, biological studies 148-03-8,  $\beta$ -Tocopherol  
328-50-7,  $\alpha$ -Ketoglutaric acid 373-49-9, Palmitoleic acid  
432-70-2,  $\alpha$ -Carotene 463-40-1, Linolenic acid 472-92-4,  
 $\delta$ -Carotene 472-93-5,  $\gamma$ -Carotene 506-12-7,  
Margaric acid 506-30-9, Arachidic acid 544-63-8, Tetradecanoic acid,  
biological studies 544-64-9, Myristoleic acid 552-94-3, Salsalate  
600-22-6, Methyl pyruvate 1002-84-2, Pentadecanoic acid 1247-42-3,  
Methyl prednisone 1406-18-4, Vitamin E 1981-50-6, Margaroleic acid  
2922-61-4, Lithium pyruvate 3385-03-3, Flunisolide 4151-33-1,  
Potassium pyruvate 5466-77-3, Ethylhexyl p-methoxycinnamate 5534-09-8,  
Beclomethasone dipropionate 6197-30-4, Octocrylene 6385-02-0,  
Meclofenamate sodium 6829-55-6, Tocotrienol 6969-49-9, Octyl  
salicylate 7235-40-7,  $\beta$ -Carotene 7439-96-5D, Manganese, pyruvate  
complexes 7616-22-0,  $\gamma$ -Tocopherol 10504-35-5,  
D-Ascorbic acid 11103-57-4, Vitamin A 13463-67-7, Titania, biological  
studies 15307-86-5, Diclofenac 15687-27-1, Ibuprofen 17407-37-3,  
Vitamin E succinate 18983-79-4, Magnesium pyruvate 21245-02-3,  
Padimate o 22071-15-4, Ketoprofen 22204-53-1, Naproxen 22494-42-4,  
Diflunisal 29204-02-2, Gadoleic acid 34597-40-5, Fenoprofen calcium  
36322-90-4, Piroxicam 38194-50-2, Sulindac 41340-25-4, Etodolac  
42924-53-8, Nabumetone 52009-14-0, Calcium pyruvate 58817-05-3  
64425-90-7, Choline magnesium trisalicylate, biological studies  
71276-50-1 74103-07-4, Ketorolac tromethamine 96436-87-2, Octyl  
p-methoxycinnamate 149732-45-6, Propanoic acid, 2-oxo-, zinc salt  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(sunscreen-wound healing compns. for treatment of sunburn)


IT 113-24-6, Sodium pyruvate

RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(sunscreen-wound healing compns. for treatment of sunburn)

RN 113-24-6 HCAPLUS

CN Propanoic acid, 2-oxo-, sodium salt (9CI) (CA INDEX NAME)



● Na

L47 ANSWER 12 OF 25 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1996:425310 HCAPLUS

DOCUMENT NUMBER: 125:67854

TITLE: Razor cartridges comprising wound healing compositions

INVENTOR(S): Martin, Alain; Vreeland, William Elbert; Booth, Anthony R.

PATENT ASSIGNEE(S): Warner-Lambert Company, USA

SOURCE: PCT Int. Appl., 117 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9610474	A1	19960411	WO 1995-US8433	19950707
W: AU, BR, CA, CN, JP, KR, MX, RU				
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
AU 9529607	A1	19960426	AU 1995-29607	19950707
EP 783398	A1	19970716	EP 1995-925499	19950707
EP 783398	B1	20020109		
R: DE, FR, GB				
JP 2002514937	T2	20020521	JP 1996-511718	19950707
PRIORITY APPLN. INFO.:			US 1994-315734	A 19940930
			US 1995-446989	A 19950522
			WO 1995-US8433	W 19950707

AB This invention pertains to therapeutic wound healing compns. useful for preventing and reducing injury to mammalian cells affixed to razor cartridges to form therapeutic razor cartridges with wound healing composition. In one embodiment of this invention the therapeutic wound healing composition comprises (a) pyruvate; (b) an antioxidant; and (c) a mixture of saturated and unsatd. fatty acids. This invention also pertains to methods for making and using the razor cartridges comprising therapeutic wound healing compns.

IC ICM B26B021-44

ICS A61K031-20

CC 63-7 (Pharmaceuticals)

Section cross-reference(s): 62

IT Anesthetics

Antihistaminics

Antioxidants

Bactericides, Disinfectants, and Antiseptics

Encapsulation

Fungicides and Fungistats

Immunostimulants

Inflammation inhibitors

Nutrients


Sunscreens

Virucides and Virustats

Wound healing promoters

(razor cartridges with wound healing compns. containing antioxidant, fatty acids, and pyruvate)

- IT 50-21-5, Lactic acid, biological studies 50-81-7, Vitamin C, biological studies 57-10-3, Hexadecanoic acid, biological studies 57-11-4, Octadecanoic acid, biological studies 58-95-7, Vitamin E acetate 59-02-9,  $\alpha$ -Tocopherol 60-33-3, 9,12-Octadecadienoic acid (Z,Z)-, biological studies 68-26-8, Retinol 79-41-4D, esters, copolymers 79-80-1, 3,4-Didehydroretinol 112-80-1, 9-Octadecenoic acid (Z)-, biological studies 113-24-6, Sodium pyruvate 127-17-3, Pyruvic acid, biological studies 127-17-3D, Pyruvic acid, esters and salts 143-07-7, Dodecanoic acid, biological studies 148-03-8,  $\beta$ -Tocopherol 328-50-7,  $\alpha$ -Ketoglutaric acid 373-49-9, Palmitoleic acid 432-70-2,  $\alpha$ -Carotene 463-40-1, Linolenic acid 472-92-4,  $\delta$ -Carotene 472-93-5,  $\gamma$ -Carotene 506-12-7, Margaric acid 506-30-9, Arachidic acid 544-63-8, Tetradecanoic acid, biological studies 544-64-9, Myristoleic acid 600-22-6, Methyl pyruvate 1002-84-2, Pentadecanoic acid 1406-18-4, Vitamin E 1406-18-4D, Vitamin E, esters and salts 1981-50-6, Margaroleic acid 2922-61-4, Lithium pyruvate 4151-33-1, Potassium pyruvate 6829-55-6, Tocotrienol 7235-40-7,  $\beta$ -Carotene 7559-04-8,  $\alpha$ -Tocoquinone 7616-22-0,  $\gamma$ -Tocopherol 9002-93-1, Triton x-100 9003-53-6, Polystyrene 10504-35-5, D-Ascorbic acid 11103-57-4, Vitamin A 17407-37-3, Vitamin E succinate 18983-79-4, Magnesium pyruvate 29204-02-2, Gadoleic acid 52009-14-0, Calcium pyruvate 61181-29-1 71276-50-1 81686-75-1 149732-45-6, Propanoic acid, 2-oxo-, zinc salt
- RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(razor cartridges with wound healing compns. containing antioxidant, fatty acids, and pyruvate)
- IT 113-24-6, Sodium pyruvate
- RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(razor cartridges with wound healing compns. containing antioxidant, fatty acids, and pyruvate)
- RN 113-24-6 HCAPLUS
- CN Propanoic acid, 2-oxo-, sodium salt (9CI) (CA INDEX NAME)



● Na

L47 ANSWER 13 OF 25 HCAPLUS COPYRIGHT 2006 ACS on STN  
 ACCESSION NUMBER: 1996:367739 HCAPLUS  
 DOCUMENT NUMBER: 125:19043  
 TITLE: Bioadhesive-wound healing composition  
 INVENTOR(S): Leung, Sau-Hung S.; Martin, Alain

PATENT ASSIGNEE(S): Warner-Lambert Company, USA  
 SOURCE: PCT Int. Appl., 159 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 28  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9606640	A1	19960307	WO 1995-US8568	19950707
W: AU, CA, JP, MX, NZ, SG				
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
US 5658956	A	19970819	US 1995-445824	19950522
AU 9530045	A1	19960322	AU 1995-30045	19950707
AU 707353	B2	19990708		
EP 779820	A1	19970625	EP 1995-926209	19950707
R: BE, CH, DE, DK, ES, FR, GB, GR, IT, LI				
JP 10505057	T2	19980519	JP 1996-508729	19950707
ZA 9507245	A	19970630	ZA 1995-7245	19950829
PRIORITY APPLN. INFO.:				
			US 1994-298521	A 19940830
			US 1995-445824	A 19950522
			US 1991-663500	B1 19910301
			US 1993-53922	B2 19930426
			WO 1995-US8568	W 19950707

AB The present invention pertains to therapeutic bioadhesive-wound healing compns. useful for treating wounds and increasing the proliferation and resuscitation rate of mammalian cells. The compns. comprise a bioadhesive agent and a therapeutically effective amount of a wound healing composition. In one embodiment the wound healing composition comprises (a) pyruvate; (b) an antioxidant; and (c) a mixture of saturated and unsatd. fatty acids. The therapeutic bioadhesive-wound healing compns. may further comprise medicaments such as antiviral agents, antikeratolytic agents, anti-inflammatory agents, antifungal agents, antibacterial agents, immunostimulating agents, and the like. The bioadhesive-wound healing compns. may be utilized in a wide variety of pharmaceutical products. This invention also relates to methods for preparing and using the bioadhesive-wound healing compns. and the pharmaceutical products in which the compns. may be used.

IC ICM A61K045-06

ICS A61K031-355

ICI A61K031-355, A61K031-20, A61K031-19

CC 63-6 (Pharmaceuticals)

IT Anesthetics

Antibiotics

Antihistaminics

Antioxidants

Bactericides, Disinfectants, and Antiseptics

Cell proliferation

Cytotoxic agents

Fungicides and Fungistats

Immunostimulants

Inflammation inhibitors

Nutrients

Sunscreens

Virucides and Virustats

Wound healing


Wound healing promoters

(bioadhesive, topical wound healing compns. containing pyruvates,

antioxidants, and fatty acids)

IT 50-02-2, Dexamethasone 50-21-5, Lactic acid, biological studies  
 50-23-7, Hydrocortisone 50-24-8, Prednisolone 50-78-2, Aspirin  
 50-81-7, Vitamin C, biological studies 53-03-2, Prednisone 53-06-5,  
 Cortisone 53-86-1, Indomethacin 56-75-7, Chloramphenicol 57-10-3,  
 Hexadecanoic acid, biological studies 57-11-4, Octadecanoic acid,  
 biological studies 57-13-6, Urea, biological studies 57-62-5,  
 Chlortetracycline 57-92-1, Streptomycin, biological studies 58-95-7,  
 Vitamin E acetate 59-01-8, Kanamycin 59-02-9,  $\alpha$ -Tocopherol  
 59-87-0, Nitrofurazone 60-33-3, 9,12-Octadecadienoic acid (Z,Z)-,  
 biological studies 60-54-8, Tetracycline 61-33-6, Penicillin G,  
 biological studies 61-68-7, Mefenamic acid 65-85-0, Benzoic acid,  
 biological studies 67-20-9, Nitrofurantoin 67-45-8, Furazolidone  
 68-26-8, Retinol 69-53-4, Ampicillin 69-72-7, biological studies  
 76-25-5, Triamcinolone acetonide 79-57-2, Oxytetracycline 79-80-1,  
 3,4-Didehydroretinol 83-43-2, Methyl prednisolone 87-08-1, Penicillin  
 V 89-57-6, Mesalamine 99-26-3, Bismuth subgallate 108-95-2, Phenol,  
 biological studies 110-44-1, Sorbic acid 112-80-1, 9-Octadecenoic acid  
 (Z)-, biological studies 113-24-6, Sodium pyruvate 114-07-8,  
 Erythromycin 118-60-5, 2-Ethylhexyl salicylate 119-13-1,  
 $\delta$ -Tocopherol 124-94-7, Triamcinolone 127-17-3, Pyruvic acid,  
 biological studies 131-57-7, Oxybenzone 134-09-8, Menthyl anthranilate  
 143-07-7, Dodecanoic acid, biological studies 147-24-0, Diphenhydramine  
 hydrochloride 148-03-8,  $\beta$ -Tocopherol 153-61-7, Cephalothin  
 302-79-4, Tretinoin 328-50-7,  $\alpha$ -Ketoglutaric acid 373-49-9,  
 Palmitoleic acid 432-70-2,  $\alpha$ -Carotene 443-48-1, Metronidazole  
 463-40-1, Linolenic acid 472-92-4,  $\delta$ -Carotene 472-93-5,  
 $\gamma$ -Carotene 506-12-7, Margaric acid 506-30-9, Arachidic  
 acid 544-63-8, Tetradecanoic acid, biological studies 544-64-9,  
 Myristoleic acid 552-94-3, Salsalate 564-25-0, Doxycycline 600-22-6,  
 Methyl pyruvate 637-58-1, Pramoxine hydrochloride 665-66-7, Amantadine  
 hydrochloride 1002-84-2, Pentadecanoic acid 1344-85-0, Bismuth  
 aluminate 1403-66-3, Gentamycin 1404-04-2, Neomycin 1405-87-4,  
 Bacitracin 1406-05-9, Penicillin 1406-11-7, Polymyxin 1406-18-4,  
 Vitamin E 1406-18-4D, Vitamin E, esters and salts 1981-50-6,  
 Margaroleic acid 2134-78-3 2922-61-4, Lithium pyruvate 3385-03-3,  
 Flunisolid 4151-33-1, Potassium pyruvate 5466-77-3, Ethylhexyl  
 p-methoxycinnamate 5534-09-8, Beclomethasone dipropionate 5536-17-4,  
 Vidarabine 5593-20-4, Betamethasone dipropionate 6197-30-4,  
 Octocrylene 6385-02-0, Meclofenamate sodium 6506-37-2, Nimorazole  
 6829-55-6, Tocotrienol 6969-49-9, Octyl salicylate 6998-60-3,  
 Rifamycin 7235-40-7,  $\beta$ -Carotene 7616-22-0,  $\gamma$ -  
 -Tocopherol 9000-30-0, Guar gum 9003-01-4, Polyacrylic acid  
 9003-97-8, Polycarbophil 9004-32-4, Sodium CM-cellulose 9004-67-5,  
 Methyl cellulose 10504-35-5, D-Ascorbic acid 11103-57-4, Vitamin A  
 11111-12-9D, Cephalosporin, derivs. 13463-67-7, Titania, biological  
 studies 14882-18-9, Bismuth subsalicylate 15307-86-5, Diclofenac  
 15686-71-2, Cephalixin 15687-27-1, Ibuprofen 17407-37-3, Vitamin E  
 succinate 18323-44-9, Clindamycin 18983-79-4, Magnesium pyruvate  
 19387-91-8, Tinidazole 21245-02-3, Padimate o 22071-15-4, Ketoprofen  
 22204-53-1, Naproxen 22494-42-4, Diflunisal 22916-47-8, Miconazole  
 23593-75-1, Clotrimazole 25655-41-8, Povidone iodine 26787-78-0,  
 Amoxicillin 29204-02-2, Gadoleic acid 30516-87-1, Zidovudine  
 34597-40-5, Fenoprofen calcium 36322-90-4, Piroxicam 36791-04-5,  
 Ribavirin 38194-50-2, Sulindac 41340-25-4, Etodolac 42924-53-8,  
 Nabumetone 52009-14-0, Calcium pyruvate 57644-54-9, Bismuth subcitrate  
 58817-05-3 59277-89-3, Acyclovir 63585-09-1, Foscarnet sodium  
 64425-90-7, Choline magnesium trisalicylate, biological studies  
 64872-76-0, Butoconazole 65899-73-2, Tioconazole 71276-50-1

74103-07-4, Ketorolac tromethamine 81686-75-1 96436-87-2, Octyl  
 p-methoxycinnamate 107910-75-8, Ganciclovir sodium 149732-45-6,  
 Propanoic acid, 2-oxo-, zinc salt 152521-52-3, Betafectin  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (bioadhesive, topical wound healing compns. containing pyruvates,  
 antioxidants, and fatty acids)  
 IT 113-24-6, Sodium pyruvate  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (bioadhesive, topical wound healing compns. containing pyruvates,  
 antioxidants, and fatty acids)  
 RN 113-24-6 HCAPLUS  
 CN Propanoic acid, 2-oxo-, sodium salt (9CI) (CA INDEX NAME)



● Na

L47 ANSWER 14 OF 25 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1996:318495 HCAPLUS

DOCUMENT NUMBER: 124:352761

TITLE: Antifungal-wound healing compositions containing  
 pyruvates and antioxidants and fatty acids

INVENTOR(S): Martin, Alain

PATENT ASSIGNEE(S): Warner-Lambert Company, USA

SOURCE: PCT Int. Appl., 114 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 28

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9603149	A1	19960208	WO 1995-US8551	19950707
W: AU, CA, JP, MX, NZ, SG				
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
US 5663208	A	19970902	US 1995-445831	19950522
AU 9530042	A1	19960222	AU 1995-30042	19950707
AU 701179	B2	19990121		
EP 773795	A1	19970521	EP 1995-926203	19950707
R: BE, CH, DE, DK, ES, FR, GB, GR, IT, LI				
JP 10503200	T2	19980324	JP 1995-505755	19950707
ZA 9506117	A	19970421	ZA 1995-6117	19950721
PRIORITY APPLN. INFO.:			US 1994-279462	A 19940722
			US 1995-445831	A 19950522
			US 1991-663500	B1 19910301
			US 1993-53922	B2 19930426
			WO 1995-US8551	W 19950707

AB Therapeutic antifungal-wound healing compns. comprise (a) pyruvate; (b) an antioxidant; and (c) a mixture of saturated and unsatd. fatty acids. The therapeutic antifungal-wound healing compns. may be utilized in a wide variety of topical and oral pharmaceutical products. A wound healing

composition contained sodium pyruvate 2, vitamin E 1, chicken fat 2, LYCD 2400U, shark liver oil 3, petrolatum 64, mineral oil 22.53, paraffin 5, and emulsifier 0.2%. The above composition was applied on a 3 cm full thickness longitudinal incision on the back of hairless mice once/day for 7 days. The composition was significantly better than preparation H and there

was

less scar tissue present at day 7 on the skin.

IC ICM A61K045-06

ICS A61K031-355

ICI A61K031-355, A61K031-20, A61K031-19

CC 63-6 (Pharmaceuticals)

IT Anesthetics

Antihistaminics

Antioxidants

Bactericides, Disinfectants, and Antiseptics

Culture media

Fungicides and Fungistats

Immunostimulants

Inflammation inhibitors

Sunscreens

Virucides and Virustats

Wound healing

(antifungal-wound healing compns. containing pyruvates and antioxidants and fatty acids)

IT 50-02-2, Dexamethasone 50-21-5, Lactic acid, biological studies  
50-23-7, Hydrocortisone 50-24-8, Prednisolone 50-78-2, Acetylsalicylic acid 50-81-7, Vitamin c, biological studies 53-03-2, Prednisone 53-06-5, Cortisone 53-86-1, Indomethacin 58-95-7, Vitamin e acetate 59-02-9,  $\alpha$ -Tocopherol 61-68-7, Mefenamic acid 76-25-5, Triamcinolone acetate 79-80-1, 3,4-Didehydroretinol 83-43-2, Methyl prednisolone 89-57-6, Mesalamine 110-44-1, Sorbic acid 113-24-6, Sodium pyruvate 119-13-1,  $\delta$ -Tocopherol 124-94-7, Triamcinolone 127-17-3, Pyruvic acid, biological studies 148-03-8,  $\beta$ -Tocopherol 328-50-7,  $\alpha$ -Ketoglutaric acid 472-92-4,  $\delta$ -Carotene 472-93-5,  $\gamma$ -Carotene 552-94-3, Salsalate 600-22-6, Methyl pyruvate 1406-18-4, Vitamin e 2922-61-4, Lithium pyruvate 3385-03-3, Flunisolide 4151-33-1, Potassium pyruvate 5534-09-8, Beclomethasone dipropionate 5593-20-4, Betamethasone dipropionate 6385-02-0, Meclofenamate sodium 6829-55-6, Tocotrienol 7235-40-7,  $\beta$ -Carotene 7488-99-5,  $\alpha$ -Carotene 7559-04-8 7616-22-0,  $\gamma$ -Tocopherol 11103-57-4, Vitamin a 15307-86-5, Diclofenac 15687-27-1, Ibuprofen 18983-79-4, Magnesium pyruvate 22071-15-4, Ketoprofen 22204-53-1, Naproxen 22494-42-4, Diflunisal 34597-40-5, Fenoprofen calcium 36322-90-4, Piroxicam 37311-39-0, Vitamin e succinate 38194-50-2, Sulindac 41340-25-4, Etodolac 42924-53-8, Nabumetone 52009-14-0, Calcium pyruvate 64425-90-7, Choline magnesium trisalicylate, biological studies 71276-50-1 74103-07-4, Ketorolac tromethamine 81686-75-1 149732-45-6  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)


(antifungal-wound healing compns. containing pyruvates and antioxidants and fatty acids)

IT 113-24-6, Sodium pyruvate

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(antifungal-wound healing compns. containing pyruvates and antioxidants and fatty acids)

RN 113-24-6 HCAPLUS  
 CN Propanoic acid, 2-oxo-, sodium salt (9CI) (CA INDEX NAME)



● Na

L47 ANSWER 15 OF 25 HCAPLUS COPYRIGHT 2006 ACS on STN  
 ACCESSION NUMBER: 1996:171907 HCAPLUS  
 DOCUMENT NUMBER: 124:212140  
 TITLE: Anti-inflammatory wound healing compositions  
 containing pyruvates and antioxidants and fatty acids  
 INVENTOR(S): Martin, Alain  
 PATENT ASSIGNEE(S): Warner-Lambert Co., USA  
 SOURCE: PCT Int. Appl., 113 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 28  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9600584	A1	19960111	WO 1995-US7942	19950622
W: AU, CA, JP, MX, NZ, SG				
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
US 5648380	A	19970715	US 1995-445845	19950522
AU 9529080	A1	19960125	AU 1995-29080	19950622
AU 701454	B2	19990128		
EP 759783	A1	19970305	EP 1995-924660	19950622
R: BE, CH, DE, DK, ES, FR, GB, GR, IT, LI				
JP 10502345	T2	19980303	JP 1995-503323	19950622
NZ 289287	A	20010223	NZ 1995-289287	19950622
ZA 9505408	A	19970401	ZA 1995-5408	19950629
PRIORITY APPLN. INFO.:			US 1994-268429	A 19940630
			US 1995-445845	A 19950522
			US 1991-663500	B1 19910301
			US 1993-53922	B2 19930426
			WO 1995-US7942	W 19950622

AB Therapeutic anti-inflammatory wound healing comps. comprise a therapeutically effective amount of one or more anti-inflammatory agents and a wound healing composition A wound healing composition contained sodium pyruvate 2

(I), vitamin E (II) 1, chicken fat 2 (III), shark liver oil 3, petrolatum 64, mineral oil 22.53, paraffin 5, emulsifier 0.2% and live yeast cell derivative 2400 U. The composition was significantly better wound healing composition than controls with no I, II, and III in healing incision wound in mice skin.

IC ICM A61K045-06

CC 63-7 (Pharmaceuticals)

Section cross-reference(s): 1




IT Acne  
 Anesthetics  
 Antihistaminics  
 Antioxidants  
 Bactericides, Disinfectants, and Antiseptics  
 Burn  
 Fungicides and Fungistats  
 Immunostimulants  
 Inflammation inhibitors  
 Nutrients  
 Sunburn and Suntan  
 Sunscreens  
 Virucides and Virustats  
 Wound healing  
     (anti-inflammatory wound healing compns. containing pyruvates and  
     antioxidants and fatty acids)

IT 50-02-2, Dexamethasone 50-23-7, Hydrocortisone 50-24-8, Prednisolone  
 50-78-2, Acetylsalicylic acid 50-81-7, Vitamin c, biological studies  
 53-03-2, Prednisone 53-06-5, Cortisone 53-86-1, Indomethacin  
 57-10-3, Palmitic acid, biological studies 57-11-4, Stearic acid,  
 biological studies 58-95-7, Vitamin e acetate 59-02-9,  
 $\alpha$ -Tocopherol 60-33-3, Linoleic acid, biological studies 61-68-7,  
 Mefenamic acid 68-26-8, Vitamin a 76-25-5, Triamcinolone acetone  
 79-80-1, 3,4-Didehydroretinol 83-43-2, Methyl prednisolone 89-57-6,  
 Mesalamine 112-80-1, Oleic acid, biological studies 113-24-6,  
 Sodium pyruvate 119-13-1,  $\delta$ -Tocopherol 124-94-7, Triamcinolone  
 127-17-3, Pyruvic acid, biological studies 143-07-7, Lauric acid,  
 biological studies 148-03-8,  $\beta$ -Tocopherol 328-50-7,  
 $\alpha$ -Ketoglutaric acid 373-49-9, Palmitoleic acid 432-70-2,  
 $\alpha$ -Carotene 472-92-4,  $\delta$ -Carotene 472-93-5,  $\gamma$ -  
 $\gamma$ -Carotene 506-12-7, Margaric acid 506-30-9, Arachidic acid 544-63-8,  
 Myristic acid, biological studies 544-64-9, Myristoleic acid 552-94-3,  
 Salicylsalicylic acid 600-22-6, Methyl pyruvate 1002-84-2,  
 Pentadecanoic acid 1406-18-4, Vitamin e 1981-50-6, Margaroleic acid  
 2922-61-4, Lithium pyruvate 3385-03-3, Flunisolide 4151-33-1,  
 Potassium pyruvate 5534-09-8, Beclomethasone dipropionate 5593-20-4,  
 Betamethasone dipropionate 6385-02-0, Meclofenamate sodium 6829-55-6,  
 Tocotrienol 7235-40-7,  $\beta$ -Carotene 7616-22-0,  $\gamma$ -  
 $\gamma$ -Tocopherol 10504-35-5, D-Ascorbic acid 15307-86-5, Diclofenac  
 15687-27-1, Ibuprofen 18983-79-4, Magnesium pyruvate 22071-15-4,  
 Ketoprofen 22204-53-1, Naproxen 22494-42-4, Diflunisal 24887-16-9,  
 Zinc, bis(2-oxopropanoato-O1,O2)-, (T-4)- 29204-02-2, Gadoleic acid  
 34597-40-5 36322-90-4, Piroxicam 37311-39-0, Vitamin e succinate  
 38194-50-2, Sulindac 41340-25-4, Etodolac 42924-53-8, Nabumetone  
 52009-14-0, Calcium pyruvate 64425-90-7, Choline magnesium  
 trisalicylate, biological studies 71276-50-1, 2H-1-Benzopyran-6-ol,  
 3,4-dihydro-2,5,7,8-tetramethyl-2-(4,8,12-trimethyltridecyl)-,  
 dihydrogenphosphate, [2R-[2R\*(4R\*,8R\*)]]- 74103-07-4, Ketorolac  
 tromethamine 145482-34-4, Manganese, bis(2-oxopropanoato-O1,O2)-  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological  
 study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES  
 (Uses)  
     (anti-inflammatory wound healing compns. containing pyruvates and  
     antioxidants and fatty acids)

IT 113-24-6, Sodium pyruvate  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological  
 study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES  
 (Uses)  
     (anti-inflammatory wound healing compns. containing pyruvates and

antioxidants and fatty acids)  
 RN 113-24-6 HCAPLUS  
 CN Propanoic acid, 2-oxo-, sodium salt (9CI) (CA INDEX NAME)



● Na


L47 ANSWER 16 OF 25 HCAPLUS COPYRIGHT 2006 ACS on STN  
 ACCESSION NUMBER: 1996:171900 HCAPLUS  
 DOCUMENT NUMBER: 124:212068  
 TITLE: Antikeratolytic wound healing compositions containing  
 pyruvates and antioxidants and fatty acids  
 INVENTOR(S): Martin, Alain  
 PATENT ASSIGNEE(S): Warner-Lambert Co., USA  
 SOURCE: PCT Int. Appl., 107 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 28  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9600572	A1	19960111	WO 1995-US7941	19950622
W: AU, CA, JP, MX, NZ, SG				
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
US 5641814	A	19970624	US 1995-445808	19950522
AU 9528707	A1	19960125	AU 1995-28707	19950622
AU 701301	B2	19990121		
EP 768877	A1	19970423	EP 1995-924046	19950622
R: BE, CH, DE, DK, ES, FR, GB, GR, IT				
JP 10502344	T2	19980303	JP 1995-503322	19950622
NZ 288995	A	20010223	NZ 1995-288995	19950622
ZA 9505409	A	19970401	ZA 1995-5409	19950629
PRIORITY APPLN. INFO.:			US 1994-268772	A 19940630
			US 1995-445808	A 19950522
			US 1991-663500	B2 19910301
			US 1993-53922	B1 19930426
			WO 1995-US7941	W 19950622

AB Therapeutic antikeratolytic wound healing comps. comprise a therapeutically effective amount of one or more antikeratolytic agents and a wound healing composition A wound healing composition contained sodium pyruvate 2 (I), vitamin E (II) 1, chicken fat 2 (III), shark liver oil 3, petrolatum 64, mineral oil 22.53, paraffin 5, emulsifier 0.2% and live yeast cell derivative 2400 U. The composition was significantly better wound healing composition than controls with no I, II, and III in healing incision wound in mice skin.

IC ICM A61K031-355  
 ICS A61K031-60

ICI A61K031-60, A61K031-355, A61K031-20, A61K031-19, A61K031-17  
 CC 63-6 (Pharmaceuticals)  
 Section cross-reference(s): 1  
 IT Acne  
 Anesthetics  
 Antihistaminics  
 Antioxidants  
 Bactericides, Disinfectants, and Antiseptics  
 Burn  
 Fungicides and Fungistats  
 Immunostimulants  
 Inflammation inhibitors  
 Nutrients  
 Reducing agents  
 Sunburn and Suntan  
 Sunscreens  
 Virucides and Virustats  
 Wound healing  
 (antikeratolytic wound healing compns. containing pyruvates and  
 antioxidants and fatty acids)  
 IT 50-21-5, Lactic acid, biological studies 50-81-7, Vitamin c, biological  
 studies 57-10-3, Palmitic acid, biological studies 57-11-4, Stearic  
 acid, biological studies 57-13-6, Urea, biological studies 58-95-7,  
 Vitamin e acetate 59-02-9,  $\alpha$ -Tocopherol 60-33-3, Linoleic acid,  
 biological studies 69-72-7, Salicylic acid, biological studies  
 79-80-1, 3,4-Didehydroretinol 112-80-1, Oleic acid, biological studies  
 113-24-6, Sodium pyruvate 119-13-1,  $\delta$ -Tocopherol  
 127-17-3, Pyruvic acid, biological studies 143-07-7, Lauric acid,  
 biological studies 148-03-8,  $\beta$ -Tocopherol 328-50-7,  
 $\alpha$ -Ketoglutaric acid 373-49-9, Palmitoleic acid 432-70-2,  
 $\alpha$ -Carotene 472-92-4,  $\delta$ -Carotene 472-93-5,  $\gamma$   
 -Carotene 506-12-7, Margaric acid 506-30-9, Arachidic acid 544-63-8,  
 Myristic acid, biological studies 544-64-9, Myristoleic acid 552-94-3,  
 Salicylsalicylic acid 600-22-6, Methyl pyruvate 1002-84-2,  
 Pentadecanoic acid 1406-18-4, Vitamin e 1981-50-6, Margaroleic acid  
 2922-61-4, Lithium pyruvate 4151-33-1, Potassium pyruvate 6829-55-6,  
 Tocotrienol 7235-40-7,  $\beta$ -Carotene 7616-22-0,  $\gamma$   
 -Tocopherol 10504-35-5, D-Ascorbic acid 11103-57-4, Vitamin a  
 18983-79-4, Magnesium pyruvate 24887-16-9, Zinc, bis(2-oxopropanoato-  
 01,02)-, (T-4)- 29204-02-2, Gadoleic acid 37311-39-0, Vitamin e  
 succinate 52009-14-0, Calcium pyruvate 71276-50-1,  
 2H-1-Benzopyran-6-ol, 3,4-dihydro-2,5,7,8-tetramethyl-2-(4,8,12-  
 trimethyltridecyl)-, dihydrogenphosphate, [2R-[2R\*(4R\*,8R\*)]]-  
 145482-34-4, Manganese, bis(2-oxopropanoato-01,02)-  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological  
 study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES  
 (Uses)  
 (antikeratolytic wound healing compns. containing pyruvates and  
 antioxidants and fatty acids)  
 IT 113-24-6, Sodium pyruvate  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological  
 study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES  
 (Uses)  
 (antikeratolytic wound healing compns. containing pyruvates and  
 antioxidants and fatty acids)  
 RN 113-24-6 HCAPLUS  
 CN Propanoic acid, 2-oxo-, sodium salt (9CI) (CA INDEX NAME)



● Na

L47 ANSWER 17 OF 25 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1995:896186 HCAPLUS

DOCUMENT NUMBER: 123:290406

TITLE: Preparation of complexes of matrix polymers with hydrogen peroxide and C1-4 mono- and C4-18 diperoxy-carboxylic acids in fluidized bed process

INVENTOR(S): Breitenbach, Joerg; Grabowski, Sven; Sanner, Axel

PATENT ASSIGNEE(S): BASF A.-G., Germany

SOURCE: Ger. Offen., 8 pp.

CODEN: GWXXBX

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 4344131	A1	19950629	DE 1993-4344131	19931223
CA 2179663	AA	19950629	CA 1994-2179663	19941210
WO 9517345	A2	19950629	WO 1994-EP4115	19941210
WO 9517345	A3	19950803		
W: CA, JP, US				
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
EP 735982	A1	19961009	EP 1995-904453	19941210
EP 735982	B1	19990421		
R: BE, CH, DE, ES, FR, GB, IT, LI, NL				
JP 09506877	T2	19970708	JP 1994-517143	19941210
ES 2129800	T3	19990616	ES 1995-904453	19941210
US 5753770	A	19980519	US 1996-663321	19960621
PRIORITY APPLN. INFO.:			DE 1993-4344131	A 19931223
			WO 1994-EP4115	W 19941210

AB Complexes are prepared in a fluidized bed process by combining matrix polymers (e.g., polymers of N-vinylcaprolactam, N-vinylpyrrolidone, and/or N-vinylimidazole, carbohydrates, and their mixts.) with aqueous solns. of H<sub>2</sub>O<sub>2</sub> and/or C1-4 mono- and C4-18 diperoxy-carboxylic acids. The complexes are useful as disinfectants, catalysts, bleaching agents, etc.

IC ICM C07C409-24

ICS C07C407-00; C08L039-04; C08L005-00; C08L003-00

ICA C08K005-14

CC 45-5 (Industrial Organic Chemicals, Leather, Fats, and Waxes)

Section cross-reference(s): 38

ST hydrogen peroxide polymer complex prepn;  
 peroxy-carboxylic acid polymer complex prepn; peroxide polymer complex prepn fluidized bed; peracetic acid polymer complex prepn;  
 vinylcaprolactam polymer peroxide complex prepn; vinylpyrrolidone polymer peroxide complex prepn; vinylimidazole polymer peroxide complex prepn;  
 carbohydrate peroxide complex prepn; granulation peroxide polymer fluidized bed

IT Fluidized beds and systems  
(for preparation of complexes of polymers with hydrogen peroxide and peroxycarboxylic acids)

IT Drying  
Granulation  
(in fluidized bed process for preparation of complexes of polymers with hydrogen peroxide and peroxycarboxylic acids)

IT 50-99-7, Glucose, processes 57-50-1, Sucrose, processes 99-20-7, Trehalose 7585-39-9,  $\beta$ -Cyclodextrin 9003-39-8, N-Vinylpyrrolidone polymer 9005-25-8, Starch, processes 9050-36-6, Maltodextrin 10016-20-3,  $\alpha$ -Cyclodextrin 17465-86-0,  $\gamma$ -Cyclodextrin 25189-83-7, N-Vinylcaprolactam polymer 25232-42-2, N-Vinylimidazole polymer 29297-55-0, N-Vinylimidazole-N-vinylpyrrolidone copolymer 30307-40-5, N-Vinylcaprolactam-N-vinylimidazole-N-vinylpyrrolidone copolymer 51987-20-3, N-Vinylcaprolactam-N-vinylpyrrolidone copolymer  
RL: PEP (Physical, engineering or chemical process); TEM (Technical or engineered material use); PROC (Process); USES (Uses)  
(fluidized bed process for preparation of complexes of hydrogen peroxide and peroxycarboxylic acids with)

IT 79-21-0, Peracetic acid 7722-84-1, Hydrogen peroxide, processes  
RL: PEP (Physical, engineering or chemical process); TEM (Technical or engineered material use); PROC (Process); USES (Uses)  
(fluidized bed process for preparation of complexes of polymers with)

IT 9003-39-8, N-Vinylpyrrolidone polymer  
RL: PEP (Physical, engineering or chemical process); TEM (Technical or engineered material use); PROC (Process); USES (Uses)  
(fluidized bed process for preparation of complexes of hydrogen peroxide and peroxycarboxylic acids with)


RN 9003-39-8 HCAPLUS

CN 2-Pyrrolidinone, 1-ethenyl-, homopolymer (9CI) (CA INDEX NAME)

CM 1

CRN 88-12-0

CMF C6 H9 N O



IT 7722-84-1, Hydrogen peroxide, processes  
RL: PEP (Physical, engineering or chemical process); TEM (Technical or engineered material use); PROC (Process); USES (Uses)  
(fluidized bed process for preparation of complexes of polymers with)

RN 7722-84-1 HCAPLUS

CN Hydrogen peroxide (H2O2) (9CI) (CA INDEX NAME)



HO-OH

L47 ANSWER 18 OF 25 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1994:437806 HCAPLUS  
 DOCUMENT NUMBER: 121:37806  
 TITLE: Chemical blackboards or sheets and writing inks therefor  
 INVENTOR(S): Zhao, Shanquan  
 PATENT ASSIGNEE(S): Peop. Rep. China  
 SOURCE: Faming Zhuanli Shenqing Gongkai Shuomingshu, 15 pp.  
 CODEN: CNXXEV  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Chinese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
CN 1073696	A	19930630	CN 1992-115211	19921225
CN 1053208	B	20000607		


PRIORITY APPLN. INFO.: CN 1992-115211 19921225  
 AB Title blackboards are prepared by coating compns. containing color developers (e.g., indicators, starch, and salts) and binders (e.g., starch, gelatins, or cellulose) on various substrates. Title inks contain oxidants, reducing agents, acidic solns., or basic solns. A white cloth or paper was coated with an aqueous solution containing starch and an aqueous KI solution was used as the writing ink.  
 IC ICM C09D009-00  
 CC 42-12 (Coatings, Inks, and Related Products)  
 IT 76-54-0, 2',7'-Dichlorofluorescein 76-59-5, Bromothymol blue 458-37-7, Curcumin 518-47-8D, Sodium fluorescein, tetrabromotetrachloro or tetrachlorotetraiodo derivs. 523-42-2, Quinoline blue 596-01-0, 1-Naphtholphthalein 2320-96-9, 4',5'-Dichlorofluorescein 4430-20-0 17372-87-1, Tetrabromofluorescein 147411-96-9, 10H-Phenoxazine-1,3-diol  
 RL: USES (Uses)  
 (coatings containing, on boards, writing inks for)  
 IT 9000-01-5, Gum arabic 9002-89-5, Poly(vinyl alcohol) 9003-39-8, Poly(vinyl pyrrolidone) 9004-32-4, CMC 9004-57-3, Ethyl cellulose 9079-65-6, Cholla gum  
 RL: TEM (Technical or engineered material use); USES (Uses)  
 (coatings, containing color developers, for writing boards, inks for)  
 IT 50-81-7, Ascorbic acid, uses 110-22-5, Acetyl peroxide 497-19-8, Sodium carbonate, uses 7631-90-5, Sodium bisulfite 7681-11-0, Potassium iodide, uses 7722-84-1, Hydrogen peroxide, uses 7778-54-3, Calcium hypochlorite  
 RL: USES (Uses)  
 (writing inks containing, boards for, chemical composition-coated)  
 IT 76-59-5, Bromothymol blue  
 RL: USES (Uses)  
 (coatings containing, on boards, writing inks for)  
 RN 76-59-5 HCAPLUS  
 CN Phenol, 4,4'-(1,1-dioxido-3H-2,1-benzoxathiol-3-ylidene)bis[2-bromo-3-methyl-6-(1-methylethyl)- (9CI) (CA INDEX NAME)



IT 9003-39-8, Poly(vinyl pyrrolidone)  
 RL: TEM (Technical or engineered material use); USES (Uses)  
 (coatings, containing color developers, for writing boards, inks for)  
 RN 9003-39-8 HCAPLUS  
 CN 2-Pyrrolidinone, 1-ethenyl-, homopolymer (9CI) (CA INDEX NAME)

CM 1

CRN 88-12-0  
 CMF C6 H9 N O



IT 7722-84-1, Hydrogen peroxide, uses  
 RL: USES (Uses)  
 (writing inks containing, boards for, chemical composition-coated)  
 RN 7722-84-1 HCAPLUS  
 CN Hydrogen peroxide (H2O2) (9CI) (CA INDEX NAME)

HO-OH


L47 ANSWER 19 OF 25 HCAPLUS COPYRIGHT 2006 ACS on STN  
 ACCESSION NUMBER: 1989:520318 HCAPLUS  
 DOCUMENT NUMBER: 111:120318  
 TITLE: Environmentally friendly, economical  
 sterilization and sanitation of wastes and  
 bioproducts  
 INVENTOR(S): Beise, Eckhard; Nordheim, Willy; Nordheim, Regina;

PATENT ASSIGNEE(S): Braeuniger, Siegfried; Baer, Manfred  
 Staatliches Amt fuer Atomsicherheit und Strahlenschutz  
 der DDR, Ger. Dem. Rep.  
 SOURCE: Ger. (East), 5 pp.  
 CODEN: GEXXA8  
 DOCUMENT TYPE: Patent  
 LANGUAGE: German  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DD 263449	A1	19890104	DD 1987-306223	19870820
PRIORITY APPLN. INFO.:			DD 1987-306223	19870820
AB	The method, suitable for treating a multiplicity of products and wastes, uses a combination of a biocide or biostat (or a mixture thereof) and ionizing irradiation, e.g. gamma rays or electron beams, to sterilize the material; the amount of irradiation is 5-95% of the usual amount required.			
IC	ICM A61L002-08 ICS A23L003-26; C02F001-30			
CC	60-2 (Waste Treatment and Disposal)			
ST	sterilization waste irradiation biocide			
IT	Swimming pools (disinfection of waters for, combined ionizing radiation and biocides in)			
IT	Wastewater treatment sludge (disinfection of, combination of biocide or biostat with ionizing radiation in)			
IT	Betaines RL: PROC (Process) (C12-14-alkyldimethyl, in inactivation of parainfluenza virus type 3 with gamma irradiation)			
IT	50-00-0, Formaldehyde, uses and miscellaneous 50-21-5, Lactic acid, uses and miscellaneous 56-81-5, Glycerol, uses and miscellaneous 62-54-4, Calcium acetate 64-17-5, Ethanol, uses and miscellaneous 64-18-6, Formic acid, uses and miscellaneous 64-19-7, Acetic acid, uses and miscellaneous 65-85-0, Benzoic acid, uses and miscellaneous 77-92-9, Citric acid, uses and miscellaneous 79-09-4, Propionic acid, uses and miscellaneous 87-69-4, Tartaric acid, uses and miscellaneous 94-13-3 99-76-3 100-97-0, Hexamethylene tetramine, uses and miscellaneous 110-44-1 120-47-8 137-40-6, Sodium propionate 141-53-7, Sodium formate 144-55-8, Sodium hydrogen carbonate, uses and miscellaneous 327-62-8, Potassium propionate 471-34-1, Calcium carbonate, uses and miscellaneous 497-19-8, Sodium carbonate, uses and miscellaneous 532-32-1, Sodium benzoate 544-17-2, Calcium formate 582-25-2, Potassium benzoate 584-08-7, Potassium carbonate 590-29-4, Potassium formate 1310-61-8, Potassium hydrogen sulfide 1313-82-2, Sodium sulfide, uses and miscellaneous 2090-05-3, Calcium benzoate 4075-81-4, Calcium propionate 5026-62-0 7440-22-4, Silver, uses and miscellaneous 7446-09-5, Sulfur dioxide, uses and miscellaneous 7492-55-9, Calcium sorbate 7647-14-5, Sodium chloride, uses and miscellaneous 7681-57-4, Sodium disulfite 7757-81-5, Sodium sorbate 7773-03-7, Potassium bisulfite 7778-54-3, Lime chloride 7782-99-2, Sulfurous acid, uses and miscellaneous 9003-53-6 9032-08-0 10599-90-3, Chloramide 16721-80-5, Sodium hydrogen sulfide 20548-54-3, Calcium sulfide 21146-90-7 24634-61-5, Potassium sorbate 35285-68-8 35285-69-9 RL: BIOL (Biological study)			



(as biocide or biostat, in combination with ionizing radiation)  
 IT 7681-57-4, Sodium disulfite  
 RL: BIOL (Biological study)  
 (as biocide or biostat, in combination with ionizing radiation)  
 RN 7681-57-4 HCAPLUS  
 CN Disulfurous acid, disodium salt (9CI) (CA INDEX NAME)



●2 Na

L47 ANSWER 20 OF 25 HCAPLUS COPYRIGHT 2006 ACS on STN  
 ACCESSION NUMBER: 1986:116070 HCAPLUS  
 DOCUMENT NUMBER: 104:116070  
 TITLE: Radiation sterilization and pasteurization  
 of solutions of some drugs and components of injection  
 solutions  
 AUTHOR(S): Safarov, S. A.  
 CORPORATE SOURCE: Inst. Biofiz., Moscow, USSR  
 SOURCE: Khimiko-Farmatsevticheskii Zhurnal (1985), 19(12),  
 1472-8  
 CODEN: KHFZAN; ISSN: 0023-1134  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Russian  
 AB The use of .gamma.-irradiation for sterilization was  
 studied by using 30 common drugs. The content and physicochem. properties  
 of drugs packaged in glass or polyethylene [9002-88-4] bottles remained  
 unchanged after .gamma.-irradiation of 25 KGy (1.8 Gy/s at room  
 temperature). Thus, .gamma.-irradiation of this dose may be used as  
 method for sterilization of pharmaceuticals, especially injections.  
 CC 63-8 (Pharmaceuticals)  
 ST gamma ray pharmaceutical sterilization  
 IT Bottles  
 (glass and polyethylene, γ -ray sterilization  
 of pharmaceutical injections in)  
 IT Gamma ray, biological effects  
 (sterilization of pharmaceuticals by)  
 IT Pharmaceuticals  
 (sterilization of, by γ -ray)  
 IT Sterilization and Disinfection  
 (γ -ray, of pharmaceutical injections)  
 IT Pharmaceuticals  
 (injections, sterilization of, by γ -ray)  
 IT 9002-88-4  
 RL: USES (Uses)  
 (bottles, γ -ray sterilization of  
 pharmaceutical injections in)  
 IT 50-99-7, biological studies 51-05-8 54-21-7 57-08-9 60-32-2  
 60-93-5 62-33-9 64-17-5, biological studies 67-48-1 76-22-2  
 98-92-0 100-97-0, biological studies 144-55-8, biological studies  
 150-59-4 299-28-5 522-40-7 614-39-1 630-56-8 7447-40-7,  
 biological studies 7460-14-2 7487-88-9, biological studies 7631-89-2

7647-14-5, biological studies 7772-98-7 9005-49-6, biological studies 10043-52-4, biological studies 12111-24-9 17224-46-3 19238-49-4 27236-88-0


RL: BIOL (Biological study)  
(injections, sterilization of, by  $\gamma$  -ray)

IT 7772-98-7

RL: BIOL (Biological study)  
(injections, sterilization of, by  $\gamma$  -ray)

RN 7772-98-7 HCAPLUS

CN Thiosulfuric acid (H<sub>2</sub>S<sub>2</sub>O<sub>3</sub>), disodium salt (9CI) (CA INDEX NAME)



●2 Na

L47 ANSWER 21 OF 25 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1985:32180 HCAPLUS

DOCUMENT NUMBER: 102:32180

TITLE: Production of polymer films containing oil-soluble materials

AUTHOR(S): Chukhadzhyan, G. A.; Sarkisyan, F. A.; Karapetyan, S. A.; Gabrielyan, E. S.

CORPORATE SOURCE: Erevan. Med. Inst., Yerevan, USSR

SOURCE: Armyanskii Khimicheskii Zhurnal (1984), 37(8), 512-17  
CODEN: AYKZAN; ISSN: 0515-9628

DOCUMENT TYPE: Journal

LANGUAGE: Russian

AB High-viscosity stable emulsions of lipophilic substances (sea buckthorn oil, propolis, cerebrosides, and lipophilic vitamins) were prepared with emulsifiers (e.g., Tween 20 [9005-64-5], Tween 80 [9005-65-6], Triton X-100 [9002-93-1], Na lauryl sulfate [151-21-3] and glycerol monostearate [31566-31-1]), preservatives (e.g., Na metabisulfite, Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>, or Me p-hydroxybenzoate [99-76-3]) hydrophilic polymers such as poly(vinylpyrrolidone) [9003-39-8], polyethylene glycol [25322-68-3] or partially hydrolyzed poly(vinyl acetate). The emulsions were used for the preparation of bilayer tissue-adhesive films. The films were wrapped and sterilized by 60Co .gamma.-radiation or UV radiation.

CC 63-7 (Pharmaceuticals)

IT 59-02-9 59-43-8, biological studies 77-92-9, biological studies

83-88-5, biological studies 99-76-3 1406-18-4 7681-38-1

7681-57-4 7757-82-6, biological studies 7772-98-7

8059-24-3 9003-20-7D, hydrolyzed 9003-39-8 11103-57-4 13870-29-6 25322-68-3

RL: BIOL (Biological study)

(emulsions containing surfactants and, for tissue-adhesive bilayer films)


IT 7681-57-4 7772-98-7

RL: BIOL (Biological study)

(emulsions containing surfactants and, for tissue-adhesive bilayer films)


RN 7681-57-4 HCAPLUS

CN Disulfurous acid, disodium salt (9CI) (CA INDEX NAME)



●2 Na


RN 7772-98-7 HCAPLUS  
 CN Thiosulfuric acid (H<sub>2</sub>S<sub>2</sub>O<sub>3</sub>), disodium salt (9CI) (CA INDEX NAME)



●2 Na

L47 ANSWER 22 OF 25 HCAPLUS COPYRIGHT 2006 ACS on STN  
 ACCESSION NUMBER: 1976:470617 HCAPLUS  
 DOCUMENT NUMBER: 85:70617  
 TITLE: Radiolysis of dilute aqueous solution of sodium iodide  
 AUTHOR(S): Shubnyakova, L. P.; Kharlamov, V. T.; Pikaev, A. K.  
 CORPORATE SOURCE: Inst. Biofiz., Moscow, USSR  
 SOURCE: Khimiya Vysokikh Energii (1976), 10(1), 49-54  
 CODEN: KHVKA0; ISSN: 0023-1193  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Russian  
 AB The sterilization of an aqueous solution of <sup>131</sup>I-labeled NaI (3 + 10-4M) by .gamma.-ray radiolysis was studied. The yields of I<sub>2</sub> and IO<sub>3</sub><sup>-</sup> and the consumption of I<sup>-</sup> were measured as a function of I<sup>-</sup> concentration and the radiation dose rate. Addition of Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> at (4-6) + 10-3M completely suppresses the radiolysis of I<sup>-</sup>.  
 CC 74-1 (Radiation Chemistry, Photochemistry, and Photographic Processes)  
 Section cross-reference(s): 63  
 ST radiolysis aq sodium iodide; sterilization aq sodium iodide radiolysis  
 IT Sterilization and Disinfection  
 (of aqueous sodium iodide solns. by γ -ray irradiation in presence of sodium thiosulfate)  
 IT Radiolysis  
 (of sodium iodide in aqueous solns. in presence of sodium thiosulfate, sterilization in relation to)  
 IT Gamma ray, chemical and physical effects  
 (sterilization by, of aqueous sodium iodide solns. in presence of sodium thiosulfate)  
 IT 7772-98-7  
 RL: USES (Uses)  
 (radiolysis of aqueous sodium iodide solns. in presence of, sterilization in relation to)

IT 7681-82-5, reactions  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (radiolysis of, in aqueous solns. in presence of sodium thiosulfate,  
 sterilization in relation to)  
 IT 7772-98-7  
 RL: USES (Uses)  
 (radiolysis of aqueous sodium iodide solns. in presence of,  
 sterilization in relation to)  
 RN 7772-98-7 HCAPLUS  
 CN Thiosulfuric acid (H<sub>2</sub>S<sub>2</sub>O<sub>3</sub>), disodium salt (9CI) (CA INDEX NAME)



●2 Na


L47 ANSWER 23 OF 25 HCAPLUS COPYRIGHT 2006 ACS on STN  
 ACCESSION NUMBER: 1967:40718 HCAPLUS  
 DOCUMENT NUMBER: 66:40718  
 TITLE: Preparation of injectable solutions  
 PATENT ASSIGNEE(S): CIBA Ltd.  
 SOURCE: Neth. Appl., 7 pp.  
 CODEN: NAXXAN  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Dutch  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
NL 6602950		19660909	NL	
FR 5250			FR	
GB 1098128			GB	
PRIORITY APPLN. INFO.:			US	19650308

AB Aqueous solns. of salts of 1-alkyl-2-pyridinium aldoxime are stabilized by addition of an ascorbic acid derivative and a H<sub>2</sub>O solution of Na metabisulfite. For example, to 325 g. 1-methyl-2-pyridiniumaldoxime chloride in 500 ml. H<sub>2</sub>O was added 10 ml. PhCH<sub>2</sub>OH, 5 g. isoascorbic acid, and 4 g. Na metabisulfite, and the volume brought to 1 l. with H<sub>2</sub>O. The solution was filtered through a sterile filter and sealed in 5-ml. ampuls.

IC A61K  
 CC 63 (Pharmaceuticals)  
 IT 89-65-6, D-erythro-Hex-2-enonic acid,  $\gamma$  -lactone 89-65-6  
 7681-57-4  
 RL: BIOL (Biological study)  
 (in stabilization of 2-formyl-1-methylpyridinium chloride oxime)  
 IT 7681-57-4  
 RL: BIOL (Biological study)  
 (in stabilization of 2-formyl-1-methylpyridinium chloride oxime)  
 RN 7681-57-4 HCAPLUS

CN Disulfurous acid, disodium salt (9CI) (CA INDEX NAME)



●2 Na

L47 ANSWER 24 OF 25 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1958:45912 HCAPLUS

DOCUMENT NUMBER: 52:45912

ORIGINAL REFERENCE NO.: 52:8263i,8264a-d

TITLE: Bacteriostatic properties of .gamma  
.-mercurated alcohols and of alkyl ethers of theseAUTHOR(S): Lebedeva, M. N.; Efremova, S. A.; Kostin, V. N.;  
Levina, R. Ya.

CORPORATE SOURCE: State Univ., Moscow

SOURCE: Vestnik Moskovskogo Universiteta (1957), 12(Ser. Mat.,  
Mekh. Astron., Fiz., Khim. No. 3), 149-58  
CODEN: VMUNAE; ISSN: 0372-6320

DOCUMENT TYPE: Journal

LANGUAGE: Unavailable

AB Compds. of the type MeC(OR)MeC(R1)MeCH2HgX, where R and R1 = H, Me, or Et and X = Cl, Br, I, CN, CNS, and alkyl groups, were tested against 15 bacteria and pathogenic fungi. The most effective 10 were (R1, R, X): Me, H, OCOMe; Me, H, Br (I); Me, H, CN (II); Me, H, CNS (III); Me, Me, OCOMe (IV); Me, Me, CN (V); H, H, CN (VI); H, Me, CNS (VII); the compound Me(CH2)3CH(OH)CH2HgBr (VIII); and HgCl2. These were effective against tuberculosis bacteria, especially IV and V which retarded its growth in dilns. of 1:131,072,000 and 1:262,144,000, resp. Generally the compds. were least effective against blue-green pus bacteria and against the typhoid-dysentery type. All have fungicidal properties; especially IV, V, and VII; the latter is recommended for clinical use. The presence of protein (in 10% serum) generally reduces the effectiveness of all the compds. by a factor of 2-4 (rarely 8-32). VIII has high bacteriostatic activity, but in the presence of the protein serum decreases more sharply than the others. When the compds. were neutralized with Na2S2O3 or cysteine, their bactericidal activity was greatly reduced. All compds. were lower in toxicity than HgCl2, especially I, III, VI, VII and VIII, and I was completely harmless to white mice in the dilns. used (1:1,000-4,000). The most active against microorganisms generally were II, IV, V, and VII; for combined high activity and low toxicity the best were VII, III, and I, which are recommended for vaccine preservation. The relation of chemical structure to activity is discussed.

CC 11C (Biological Chemistry: Microbiology)

IT Bactericides, Disinfectants and Antiseptics

Fungicides or Fungistats

(alcs. (γ -mercurated) and their alkyl ethers as)


IT Tuberculosis

(antitubercular substances, γ -mercurated alcs. and  
their alkyl ethers as)

IT Proteins

(bactericidal action of γ -mercurated alcs. and their  
alkyl ethers in presence of)


IT Alcohols  
 (bactericidal effect of  $\gamma$ -mercurated, and their alkyl ethers)  
 IT Vaccines  
 (preservation with  $\gamma$ -mercurated alcs. and their alkyl ethers)  
 IT 52-90-4, Cysteine 7772-98-7, Sodium thiosulfate  
 (effect on bactericidal action of  $\gamma$ -mercurated alcs. and their alkyl ethers)  
 IT 7772-98-7, Sodium thiosulfate  
 (effect on bactericidal action of  $\gamma$ -mercurated alcs. and their alkyl ethers)  
 RN 7772-98-7 HCAPLUS  
 CN Thiosulfuric acid (H<sub>2</sub>S<sub>2</sub>O<sub>3</sub>), disodium salt (9CI) (CA INDEX NAME)



●2 Na

L47 ANSWER 25 OF 25 HCAPLUS COPYRIGHT 2006 ACS on STN  
 ACCESSION NUMBER: 1954:66042 HCAPLUS  
 DOCUMENT NUMBER: 48:66042  
 ORIGINAL REFERENCE NO.: 48:11723a-c  
 TITLE: Preparation of cod-liver residues and vitamin B12 concentrates  
 AUTHOR(S): Truscott, Beryl; Gage, D. G.; Hoogland, P. L.  
 SOURCE: Journal of the Fisheries Research Board of Canada (1954), 11, 355-61  
 CODEN: JFRBAK; ISSN: 0015-296X  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Unavailable  
 AB In a study of the vitamin B12 content of cod-liver exts., two methods were used to prepare fresh residue. In the Vandenheuvel method homogenized livers were heated at 60° and mixed with 7N NaOH to pH 8.0-8.5, the residue being brought, after centrifugation, to pH 6.0-6.5 with 4N HCl. In the 2nd method, homogenized livers were warmed in water to 85°, quickly heated to 15 lb. pressure, and cooled to 100°. The mixture was centrifuged while still hot, and the oil and the residue were collected. The Vandenheuvel method produces very good results. The fresh residue could be dried without appreciable loss of vitamin B12 activity on a double drum dryer. The most satisfactory results in defatting the residue were obtained with (CH<sub>2</sub>Cl)<sub>2</sub>. Concentrates with vitamin B12 activity equivalent to 2-4 .gamma. per ml. were produced by extraction of the dried defatted residue with H<sub>2</sub>O and evaporation in vacuo.  
 CC 17 (Pharmaceuticals, Cosmetics, and Perfumes)  
 IT 1314-13-2, Zinc oxide 7772-98-7, Sodium thiosulfate  
 (as disinfectant)  
 IT 7772-98-7, Sodium thiosulfate  
 (as disinfectant)  
 RN 7772-98-7 HCAPLUS

CN Thiosulfuric acid (H<sub>2</sub>S<sub>2</sub>O<sub>3</sub>), disodium salt (9CI) (CA INDEX NAME)



●2 Na